

# Cyanobacteria, cyanotoxins, and their histopathological effects on fish tissues in Fehérvárcsurgó reservoir, Hungary

Damjana Drobac Backović · Nada Tokodi D· Zoran Marinović · Jelena Lujić · Tamara Dulić · Snežana B. Simić · Nevena B. Đorđević · Nevena Kitanović · Ilija Šćekić · Béla Urbányi · Jussi Meriluoto · Zorica Svirčev

Received: 1 March 2021 / Accepted: 17 July 2021

© The Author(s) 2021

Abstract Cyanobacteria are important members of lake plankton, but they have the ability to form blooms and produce cyanotoxins and thus cause a number of adverse effects. Freshwater ecosystems around the world have been investigated for the distribution of cyanobacteria and their toxins and the effects they have on the ecosystems. Similar research was performed on the Fehérvárcsurgó reservoir in Hungary during 2018. Cyanobacteria were present and blooming, and the highest abundance was

Highlights

- Cyanobacterial blooms and cyanotoxins are common worldwide, thus, monitoring and data on their adverse effects are required
- Six cyanobacterial species were present and blooming in Fehérvárcsurgó reservoir during 2018, a previously unexplored site in Hungary
- Microcystin (mcyE) and the saxitoxin (sxtG) encoding genes were detected in the biomass, and microcystin-RR was found in one sample
- Microcystins were not detected in fish tissue, but histopathological alterations in gills and kidneys were observed

D. Drobac Backović · N. Tokodi (☒) · Z. Marinović · J. Meriluoto · Z. Svirčev Faculty of Sciences, Department of Biology and Ecology, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia e-mail: nada.tokodi@dbe.uns.ac.rs

## N. Tokodi

Faculty of Biochemistry, Biophysics and Biotechnology, Laboratory of Metabolomics, Jagiellonian University, Gronostajowa 7, 30387 Krakow, Poland

Published online: 06 August 2021

recorded in July (2,822,000 cells/mL). The species present were Aphanizomenon flos-aquae, Microcystis flos-aquae, Microcystis wesenbergii, Cuspidothrix issatschenkoi, Dolichospermum flos-aquae, and Snowella litoralis. In July and September, the microcystin encoding gene mcyE and the saxitoxin encoding gene sxtG were amplified in the biomass samples. While a low concentration of microcystin-RR was found in one water sample from July, analyses of Abramis brama and Carassius gibelio caught from the reservoir did not show the presence of the investigated microcystins in the fish tissue. However, several histopathological changes, predominantly in gills and kidneys, were observed in the fish, and the

Z. Marinović  $\cdot$  N. Kitanović  $\cdot$  I. Šćekić  $\cdot$  B. Urbányi Department of Aquaculture, Hungarian University of Agriculture and Life Sciences, Páter Károly u. 1, Gödöllő 2100, Hungary

## J. Lujić

Department of Biomedical Sciences, Center for Reproductive Genomics, Cornell University, Ithaca, NY, USA

T. Dulić · J. Meriluoto · Z. Svirčev Faculty of Science and Engineering, Biochemistry, Åbo Akademi University, Tykistökatu 6 A, 20520 Turku, Finland

S. B. Simić · N. B. Đorđević Faculty of Science, Department of Biology and Ecology, University of Kragujevac, Radoja Domanovića 12, 34000 Kragujevac, Serbia



damage was more severe during May and especially July, which coincides with the increase in cyanobacterial biomass during the summer months. Cyanobacteria may thus have adverse effects in this ecosystem.

**Keywords** Cyanobacterial blooming · Cyanotoxin encoding genes · Microcystin · Fish histology

#### Introduction

Cyanobacteria are ancient and ubiquitous organisms that have managed to conquer numerous and diverse ecosystems. It has been estimated that the total global cyanobacterial biomass is in the order of  $3 \times 10^{14}$  g of carbon, surpassing a thousand million metric tons ( $10^{15}$  g) of wet biomass (Garcia-Pichel et al., 2003). Consequently, toxic metabolic products of cyanobacteria — cyanotoxins — are also common, as it was corroborated in a recent review article by Svirčev et al. (2019). The review reports 1118 recorded identifications of major cyanotoxins in 869 freshwater ecosystems from 66 countries worldwide. In Europe, cyanotoxins were documented in 273 freshwater ecosystems (predominantly in lakes) within 25 countries. Microcystins (MCs) were the most commonly detected cyanotoxins (58%) in these studies (Svirčev et al., 2019).

Cyanotoxins were found to be associated with adverse health effects in humans and animals, and MCs and anatoxins (ATX) were presumed causative agents in the majority of the poisonings (Svirčev et al., 2017, 2019). Poisoning cases were recorded in freshwater ecosystems across Europe: the UK, Belgium, Denmark, Finland, France, Greece, Italy, Netherlands, Norway, Sweden, Switzerland, Spain, Serbia, and Hungary (Svirčev et al., 2019). The risks caused by cyanobacterial blooms and cyanotoxins to human health are real and possibly increasing as cyanobacteria vigorously thrive in a large number of aquatic ecosystems in many countries. The synthesis of monitoring data with toxicological, epidemiological, and health data gives solid evidence of adverse effects caused by cyanobacteria and their toxins (Svirčev et al., 2017).

The objective of the present research was to monitor the presence of cyanobacteria and cyanotoxins, as well as to collect data on their effects, in Fehérvárcsurgó reservoir in Hungary. The reservoir constitutes an important freshwater ecosystem that protects the surrounding area from flooding. Due to mining activities, and the existence of a large number of sandstones in the area, a large amount of wastewater ends up in Fehérvárcsurgó reservoir (Nagy et al., 2014). Also, wastewater from the surrounding settlements, as well as organic and inorganic substances that are washed away from the surrounding agricultural land, reaches the reservoir. In the recent decades, blooms of potentially toxic cyanobacterial species were noted, but a very limited amount of data is available (Törökné, 1999; Törökné et al., 2001). The investigation consisted of qualitative and quantitative analyses of cyanobacterial presence, detection of cyanotoxin encoding genes within the cyanobacterial biomass, detection of cyanotoxins in water and fish tissues, and histopathological studies of fish tissues. Similar studies were conducted in the neighboring country, Serbia, on commercial fishponds and lake Ludaš, by the same group of authors (Drobac et al., 2016; Tokodi et al., 2018, 2020), and recently in Hungary on Lake Balaton and Kis-Balaton Water Protection System (KBWPS) (Marinović et al., 2021). The data arising from this type of research and monitoring at the local, regional, and global scale contribute to assessing the risks and consequences cyanobacteria and cyanotoxins can have on human beings and the environment.

## Materials and methods

## Sampling site

Fehérvárcsurgó reservoir is located north of Gaja River, the largest water flow of East Bakony, a mountainous region north of Lake Balaton, Hungary. The reservoir was formed in 1971 and later used for irrigation purposes of cabbage fields. Nowadays, the reservoir is utilized for flood control, industrial and drinking water abstraction, water quality improvement, irrigation, but mostly for fishing and swimming as a popular holiday resort. Its average water depth is 6 to 8 m, but its deepest points reach up to 15 m. The maximal water surface can be up to 223 ha.

## Sampling of water and fish

The sampling of water and fish from the reservoir (47°17′10.4″N, 18°14′50.6″E) was performed during four separate sampling surveys in the spring (April and May), summer (July), and autumn (September) of 2018. Water (12 L) for qualitative and quantitative analyses of cyanobacteria was collected by sweeping with a plankton net (net frame 25 cm ø, net mesh



23 µm) at a depth of 0.3 m (Marinović et al., 2021; Tokodi et al., 2020). All samples were immediately preserved in 10% neutral-buffered formalin (NBF). About 1.5 to 2 L of reservoir water was collected for cyanotoxin analyses during the investigation in July and September of the same year and frozen until the start of the analysis. Fish sampling was carried out in the center of the reservoir with a standard electrofishing device (Marinović et al., 2021; Tokodi et al., 2020). The fish were sampled during all four sampling surveys and consisted of common bream — Abramis brama (Linnaeus 1758) and Prussian carp — Carassius gibelio (Bloch 1782) (Table 1) of which the latter was sampled only in July. The following organs were collected and pooled per species for each season for MC analysis: gills, muscles, liver, kidney, spleen, intestines, and gonads (testes and ovaries). Tissue samples were also used for histopathological analyses after they were dissected and fixed in 10% neutral-buffered formalin (NBF).

Qualitative and quantitative analyses of cyanobacteria

Cyanobacterial taxa were identified at  $100 \times, 400 \times$  magnification in accordance with several taxonomic keys (Komárek, 2013; Komárek & Anagnostidis, 1998, 2005) under the Motic BA310 light microscope equipped with a digital camera (Bresser 9 MP) and Micro Cam Lab software. The quantitative analysis of phytoplankton was made with a Motic AE 2000 inverted microscope using the Utermöhl method (Utermöhl, 1958), expressing data as number of cells per ml (Marinović et al., 2021; Tokodi et al., 2020).

## Cyanotoxin encoding gene analyses

As a part of the same project, cyanotoxin encoding gene analyses were conducted as previously described in Tokodi et al. (2020) and Marinović et al. (2021). A short description of the procedure is presented below.

**Table 1** Fish species caught in Fehérvárcsurgó reservoir during 2018

Month	Species	Number of individuals	Gender ratio (male/ female)	TL (mm)
April	Abramis brama	5	3:2	$285 \pm 58$
May	Abramis brama	7	5:2	$305 \pm 51$
July	Carassius gibelio	5	1:4	$112\pm 6$
September	Abramis brama	1	_	135

## Reference strains for PCR analysis

Reference strains were procured from Finnish Environment Institute (SYKE), Pasteur Culture Collection (PCC), Australian National Algae Culture Collection (CS), and National Institute for Environmental Studies Microbial Culture Collection (NIES): MC producer: PCC7820 (Microcystis); CYN producers: CS-506 (Cylindrospermopsis); STX producer: CS-537/13 (Dolichospermum); and ATX-a producer: Dolichospermum 123 (former Anabaena 123) (SYKE).

# Extraction of DNA

Water samples of 100-250 mL (depending on the bloom density) were filtered on 2 µm membrane filters (pore size 2–3 µm), and the filters with biomass (filtrides) were freeze-dried. The genomic DNA from both the freeze-dried filters and reference strains (10 mg samples) were extracted using the DNeasy Plant Mini Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. Minimal modifications were made to successfully extract DNA from the filtrides (double amount of Buffer AP1, Buffer P3, and RNase A was added to suspend the filters entirely). Homogenization of the filter samples was achieved using zirconia/silica disruption beads (0.5 mm) and subsequent vortexing for 1 min. The success of the extraction was examined with a spectrophotometer (NanoDrop ND-1000, Thermo Scientific). The A<sub>260</sub>/ A<sub>280</sub> ratio varied between 1.22 and 2.04, indicating an adequate amount of extracted DNA in the samples.

# Qualitative PCR

The presence of MC (mcyE), CYN (cyrJ), STX (sxtG, sxtS), and ATX (anaC) synthetase genes was assessed by qualitative PCR. The PCR reaction mixtures (20  $\mu$ L) contained 1×Phire Reaction Buffer, 0.4  $\mu$ L Phire II HotStart polymerase (Thermo Scientific), 0.2



mM dNTPs (Thermo Scientific), 0.5 µL forward and reverse primers (Table 2), 2 µL of a template, and sterile deionized water (Marinović et al., 2021; Tokodi et al., 2020). The following protocols were used to run PCRs on a C1000 Touch Thermal Cycler (Bio-Rad): initial denaturation for 30 s at 98 °C; 40 cycles of 5 s at 98 °C, 5 s at 62 °C (cyrJ\_F, cyrJ\_R, sxtG432\_F, sxtG928\_R, sxtS205\_F, sxtS566\_R), or 61 °C (HEPF, HEPR), or 52 °C (anaC-genF, anaC-genR) and 10 s at 72 °C; and a final extension of 1 min at 72 °C. The potential inhibition of PCRs was examined with an exogenous amplification control template, which contained 1 µL:1 µL reference/sample. The following strains were used in the control template: PCC7820 for mcyE, CS-506 for cyrJ, CS-537/13 for sxtG and sxtS, and Dolichospermum 123 for anaC. A 1.5% Top Vision agarose gel (Thermo Scientific) dyed with SYBR® Safe DNA gel stain was used to visualize the PCR products. The Quantity One software (v. 4.6.9) of Gel DocTM XR (Bio-Rad) was used to document the observed bands.

## Cyanotoxin analyses

In order to guarantee the comparability of the results with earlier results generated within the same project, cyanotoxin analyses in water and tissues were conducted as previously described in Tokodi et al. (2020) and Marinović et al. (2021). A short description of the procedure is provided below.

## Preparation of water samples for LC-MS/MS

Depending on the bloom density, 100-250 mL of water samples was filtered (pore size 2-3  $\mu$ m). The biomass on the filter was freeze-dried and the filtrides

placed in glass tubes. The toxin extraction was accomplished by adding 3 mL of 75% MeOH and 15 min of bath ultrasonication. Samples were then further extracted with a Bandelin Sonopuls HD 2070 micro-tip probe sonicator for 1 min (30% pulse and 30% energy). After centrifugation of the extracts for 10 min at  $10,000\times g$ , 2 mL of the supernatant was evaporated to dryness (50 °C, nitrogen flow) in glass tubes. The sample was redissolved in 200  $\mu$ L 75% MeOH and in 200  $\mu$ L H<sub>2</sub>O for the analysis of MC and CYN, respectively. After filtration of the samples (0.2  $\mu$ m GHP ACRODISC 13 PALL Corporation) into inserts, they were ready for LC–MS/MS analysis.

Partial volumes of the filtrates were concentrated by solid-phase extraction (SPE) on Waters Oasis HLB (30 mg). Elution was performed with 5 mL 90% MeOH, and samples were placed into glass tubes. Evaporation was achieved using nitrogen flow, after which samples were redissolved in 200  $\mu L$  of 75% MeOH and filtered (0.2  $\mu m$  GHP ACRODISC 13 PALL Corporation) into inserts for LC–MS/MS analysis. Both fractions of the water samples were combined and presented in results.

## Preparation of fish tissue samples for LC-MS/MS

Fish liver, gills, intestines, muscle, or whole entrails (if there was not enough material) of *C. gibelio* or *A. brama* from Fehérváricsurgó reservoir were analyzed for MCs by LC–MS/MS. Tissues were homogenized, then freeze-dried, and samples of the same organ of all the individuals of the same species were pooled together. Around 400 mg of freeze-dried fish tissue samples was placed into glass tubes, and 10 mL of 75% MeOH was added for extraction of cyanotoxins

**Table 2** List of primers used for qualitative PCR

Gene	Primer	5′–3′ sequence	Reference
mcyE	HEPF	TTTGGGGTTAACTTTTTTGGGCATAGTC	Dittman et al. (2017)
	HEPR	AATTCTTGAGGCTGTAAATCGGGTTT	
cyrJ	cyrJ_F	TTCTCTCCTTTCCCTATCTCTTTATC	Mazmouz et al. (2010)
	cyrJ_R	GCTACGGTGCTGTACCAAGGGGC	
sxtG	sxtG432_F	AATGGCAGATCGCAACCGCTAT	Savela et al. (2015)
	sxtG928_R	ACATTCAACCCTGCCCATTCACT	
sxtS	sxtS205_F	GGAGTATTDGCGGGTGACTATGA	Savela et al. (2017)
	sxtS566_R	GGTGGCTACTTGGTATAACTCGCA	
anaC	anaC-genF	TCTGGTATTCAGTCCCCTCTAT	Rantala-Ylinen et al. (2011)
	anaC-genR	CCCAATAGCCTGTCATCAA	



Environ Monit Assess (2021) 193:554 Page 5 of 14 554

overnight. Probe homogenization was performed on ice for 30 s, ultrasonication in a bath sonicator for 15 min, and further extraction with a Bandelin Sonopuls HD 2070 micro-tip probe sonicator for 1 min (30% pulse and 30% energy). After centrifugation for 10 min at  $10,000 \times g$ , 5 mL of hexane to 10 mL of the obtained supernatants was added. The hexane (lipid) layer was discarded by a glass pipette. The samples were first diluted with water and then concentrated by SPE (Waters Oasis HLB 30 mg) and eluted with 5 mL 90% MeOH. Two milliliters of the samples was evaporated (50 °C nitrogen flow) in glass tubes and then redissolved in 200 µL 25% MeOH and filtered (0.2 µm GHP ACRODISC 13 PALL Corporation) into inserts. The fish tissue samples were then ready for LC-MS/MS analysis.

#### LC-MS/MS

Toxin analyses were performed by LC-MS/MS (Marinović et al., 2021; Tokodi et al., 2020). The analytical targets consisted of nine MC variants (MC-dmRR, MC-RR, MC-dmYR, MC-YR, MC-dmLR, MC-LR, MC-LY, MC-LW and MC-LF) and CYN.

# Analyses of fish histopathology

Cyprinus carpio carp (6 individuals; 3 males/3 females; TL 320±88 mm) was bred at the Department of Aquaculture, Hungarian University of Agriculture and Life Sciences (Gödöllő, Hungary) and used as control in this study. In preparation of fish tissue samples, the similar methodology as described in Tokodi et al. (2020) and Marinović et al. (2021) was used. In short, all tissue samples collected from different organs (gills, liver, kidney, muscle, spleen, intestine, and gonads) were placed in 10% NBF immediately after dissection, and after at least 3 days of fixation, the samples were processed by standard histological procedures. In addition, gill and muscle samples were decalcified before the histological processing using a 75% RDO Rapid Decalcifier solution (Apex Engineering Products Corporation). For tissue processing, the samples were dehydrated in a graded ethanol series, cleared in xylol and subsequently embedded in paraffin wax blocks. Three 5 µm thin sections per tissue per individual were cut and placed onto glass slides and stained with standard hematoxylin and eosin (H&E) staining procedure. The sections were examined under a microscope (Nikon Eclipse 600) and photographed (QImaging Micro Publisher 3.0 digital camera).

#### Results

Presence of cyanobacterial species in water samples

Six cyanobacterial species were recorded in Fehérvárcsurgó reservoir. The most dominant was *Aphanizomenon flos-aquae* Ralfs ex Bornet & Flahault (Table 3).

Presence of cyanotoxin encoding genes in biomass samples

The MC encoding gene mcyE and the STX encoding gene sxtG were amplified in biomass samples collected in July and September, while the amplicons of sxtS were not observed. The CYN encoding gene cyrJ and the ATX-a encoding gene anaC were also not amplified in this study (Fig. 1).

Presence of cyanotoxins in water and fish samples

Cyanobacterial biomass and water samples were tested for the presence of several MC variants and CYN (CYN was analyzed only in the biomass). Only MC-RR was found at a trace concentration (<0.01  $\mu$ g/L) in the sample from July 2018. The investigated MC variants were not detected in the tissue samples originating from two fish species sampled in spring, summer, and autumn.

Histopathological alterations in fish samples

Because of the close taxonomical relation between the common bream and Prussian carp, histopathological results are shown together. Since only one individual was caught during sampling in September 2018, results for this period were not analyzed and presented. The organs of control fish raised at the Department of Aquaculture, Hungarian University of Agriculture and Life Sciences, had a normal histological structure of all the organs.

Control individuals displayed a normal cord-like parenchyma with polygonal hepatocytes and centrally position nuclei with visible nucleoli (Fig. 2A, B).



Table 3 Composition of cyanobacteria from Fehérvárcsurgó reservoir in 2018

Taxon	April	May	July	September
Cyanobacteria	cells/mL	cells/mL	cells/mL	cells/mL
Aphanizomenon flos-aquae Ralfs ex Bornet & Flahault	93,500	970,000	2,124,000	1,110,000
Cuspidothrix issatschenkoi (Usachev) Rajaniemi et al	_	75,000	287,600	189,000
Dolichospermum flos-aquae (Brébisson ex Bornet & Flahault) Wacklin, Hoffmann & Komárek	-	4,200	15,200	12,100
Microcystis flos-aquae (Wittrock) Kirchner	1,100	158,000	205,700	162,000
Microcystis wesenbergii (Komárek) Komárek ex Komárek	10,800	46,000	158,300	154,000
Snowella litoralis (Häyrén) Komárek & Hindák	_	_	31,200	11,800
$\Sigma$	165,400	1,418,600	2,822,000	1,638,900

<sup>-</sup> not present in the sample


Most individuals from the Fehérvárcsurgó reservoir displayed normal hepatic structure. The most common alteration was hepatocyte vacuolization to various extents, which was most pronounced during April (Fig. 2C, D). During May (Fig. 2E) and July (Fig. 2F), almost no histopathological changes were observed.

Kidneys of control individuals displayed normal renal histology (Fig. 3A, B). Individuals from the Fehérvárcsurgó reservoir displayed several alterations throughout the seasons. The most commonly observed alterations were vacuolization of the tubular epithelium (of both proximal and distal tubules: Fig. 3C–F), which led to occasional detachment of the tubular basal lamina (Fig. 3E). Changes within the renal corpuscles were also observed; they involved dilatations of the Bowman's capsule and mild glomerular atrophy. Lymphocyte and macrophage infiltration (Fig. 3D) was also noted. These changes were the most intense during May and July, when very intense vacuolation of the tubules was detected. However, only mild vacuolations

were observed in April. Nuclear damage in the form of karyolysis was also very intense and frequent, mainly during April but also during July (Fig. 3F).

Gills of control individuals displayed normal gill histology (Fig. 4A, B). The most frequently observed gill alterations in individuals caught from the Fehérvárcsurgó reservoir were hypertrophy of epithelium (Fig. 4C), clubbing of secondary lamellae, telangiectasia, edema and epithelial lifting, hyperemia (Fig. 4D), and hyperplasia of the lamellar epithelium and the interlamellar cell mass. All mentioned alterations were noted during April. During May, hyperplasia was significantly more pronounced and often led to complete fusions of secondary lamellae (Fig. 4E). Also, in some cases, hyperplasia occurred together with epithelial lifting (Fig. 4F). However, during July, histopathological changes were not recorded, and the gills morphologically looked like those of control fish (Fig. 4G, H).

No significant histopathological alterations were observed in the other examined organs of fish from




**Fig. 1** Agarose gel showing presence of cyanotoxin encoding genes in biomass samples, where the *mcyE* gene amplicons (**A**) and *sxtG* gene amplicons (**B**) were observed in July and September 2018 samples from Fehérvárcsurgó reservoir. Legend:

R1, R2, R3 — reference strains; S1, S2 — samples from July and September, respectively; B - blank control, C - exogenous amplification control; L - ladder



Environ Monit Assess (2021) 193:554 Page 7 of 14 554



**Fig. 2** Livers of control individuals (A, B), as well as of individuals sampled from the Fehérvárcsurgó reservoir during April (C, D), May (E), and July (F). (A, B) Controls displayed normal hepatic structure. (C, D) Vacuolization of the cyto-

plasm (microvesicular steatosis). During May (E) and July (F), no significant histopathological alterations were noted. H&E staining. A, C, E, F 50  $\mu$ m; B, D 25  $\mu$ m

the Fehérvárcsurgó reservoir. The intestines (Fig. 5B), spleen (Fig. 5D), testes (Fig. 5F), ovaries (Fig. 5H), and muscles (Fig. 5J) showed a structure similar to that of the controls (Fig. A, C, E, G, I).


## Discussion

As in every lake, the plankton community in the Fehérvárcsurgó reservoir was changing during the year.

During the first sampling in April, three cyanobacterial species were recorded: *Aphanizomenon flos-aquae*, *Microcystis wesenbergii*, and *M. flos-aquae*. The number of cells of the first two species listed exceeded 10,000 cells per mL, which is defined as blooming (Falconer, 1998). The dominant *A. flos-aquae* is one of the most common species of cyanobacteria in freshwater ecosystems. It can produce CYN and STX, while some strains also possess the microcystin synthetase (*mcy*) gene cluster; thus, the production of MCs under



554 Page 8 of 14 Environ Monit Assess (2021) 193:554



**Fig. 3** Kidneys of control individuals (**A**, **B**), as well as of individuals sampled from the Fehérvárcsurgó reservoir during April (**C**), May (**D**, **E**) and July (**F**). **A**, **B** Controls displayed a normal renal structure. During April (**C**), we observed vacuolization of tubular epithelial cells, which became much more

intense during May (**D**, **E**) and July (**F**). Detachment of the tubular basal lamina (**E**; arrow), lymphocyte infiltration (**D**; star), and karyolysis (**F**; arrow) was also observed. H&E staining. **A**, **C**, **D** 50 µm; **B**, **E**, **F** 25 µm

some physiological and environmental conditions can occur (Lyon-Colbert et al., 2018). These cyanotoxins can be detrimental to aquatic animals, birds, domestic animals, and humans (Lyon-Colbert et al., 2018). Microcystis spp. is a frequently occurring and one of the main bloom-forming genera in eutrophic freshwaters well known for the production of MCs. It is assumed that in Europe, several morphospecies (e.g., M. aeruginosa) have a considerably greater portion of MC producers than others (e.g., M. flos-aquae) (Via-Ordorika et al., 2004). Although most of the analyzed M. wesenbergii strains are non-toxic, several strains were observed to produce MCs (Otsuka et al., 1999; Stoyneva-Gärtner et al., 2021). Thus, the representatives of this species should not be generally considered non-toxic. This morphospecies has a global distribution, including Europe (Jasprica et al., 2005), where it can be found less often than in some other parts of the world such as China (Xu et al., 2008).


In May and July, three further species appeared: *Cuspidothrix issatschenkoi* (synonym: *Aphanizomenon issatschenkoi*), *Dolichospermum flos-aquae* (synonym: *Anabaena flos-aquae*), and finally *Snowella litoralis*. *C. issatschenkoi* occurs occasionally and uncommonly forms blooms in the northern temperate zone, unlike in the southern temperate zone (Ballot et al., 2018). It is common in Japan, but in Europe, it is an invasive and

expanding cyanobacterium (Hodoki et al., 2012). It has a potential to produce ATX-a, homoanatoxin-a, and STXs (as seen in Japan, China, New Zealand, Germany and Portugal), but further knowledge on the worldwide distribution of ATX-a-producing strains of C. issatschenkoi is still insubstantial (Ballot et al., 2018; Hodoki et al., 2012). Therefore, it is interesting that a blooming of this species was recorded in the investigated reservoir from May to September 2018. Strains of another observed cyanobacterium, D. flos-aquae, have been found to produce ATX-a, ATX-a(S), and MCs in Finland, Japan, Canada, and USA (Bernard et al., 2017). Finally, S. litoralis is a cyanobacterium that was not extensively explored. It seems that species belonging to the genera Snowella are habitual in lakes of central Europe and Scandinavia, usually at low biomass levels, and they compose a significant portion of cyanobacterial biomass only sporadically (Rajaniemi-Wacklin et al., 2006). They are often present in late summer and autumn (Rajaniemi-Wacklin et al., 2006), just as was the case in Fehérvárcsurgó reservoir.

As it was expected in the summer, when the conditions for the blooming of cyanobacteria were the most favorable (higher temperature, light intensity and duration, nutrient enrichment), the number of individual and total cyanobacteria was the highest, with the most dominant species still being *A. flos-aquae*.



Environ Monit Assess (2021) 193:554 Page 9 of 14 554



**Fig. 4** Gills of control individuals (**A**, **B**), as well as of individuals sampled from the Fehérvárcsurgó reservoir during April (**C**, **D**), May (**E**, **F**) and July (**G**, **H**). (**A**, **B**) Controls displayed a normal gill structure. Hypertrophy of epithelium (**C**; arrows), hyperemia (**D**, arrow), and hyperplasia of the lamellar and interlamellar epithelium (**E**, **F**) were observed. During


May (**E**, **F**), hyperplasia led to complete fusions of secondary lamellae (**D**), often observed together with epithelial lifting (**F**; arrow). During July (**G**, **H**), no histopathological alterations were noted. H&E staining. **A**, **C**, **E**, **G** 100  $\mu$ m; **B**, **D**, **F**, **H** 50  $\mu$ m

At the end of the summer, the number of cyanobacteria gradually began to decline. A higher production of cyanotoxins can be expected during favorable growth conditions, and microcystin-RR (MC-RR) was detected in July. While common among MCs, MC-RR is just one of over 279 known variants of MCs (Bouaïcha et al., 2019). Scarce previous studies on Fehérvárcsurgó reservoir have shown that a sample of *M. aeruginosa* collected from the reservoir in September 1995 contained MC-RR (0.17 mg/g), MC-LR (0.38 mg/g), and MC-YR (0.33 mg/g). The sample also caused mortality of Tamnocephalus larvae and had a minimal lethal dose of 50 mg/kg in the intraperitoneal mouse assay (Törökné, 1999), induced sensitization in 30% of albino guinea pigs, and provoked slight skin and eye irritation in albino rabbits (Törökné et al., 2001). During our research, the species M. aeruginosa was not recorded in Fehérvárcsurgó reservoir.

The potential ability of cyanobacteria in Fehérvárcsurgó reservoir to produce cyanotoxins was supported by the presence of cyanotoxin encoding genes. During July and September, the MC encoding gene mcyE and the STX encoding gene sxtG were amplified in the biomass samples. The observed amplicons of the mcyE gene indicate the presence of MC-producing cyanobacteria in Fehérvárcsurgó reservoir, which is in accordance with the species observed and the obtained LC-MS/MS results. Amplification of the sxtG gene was observed in both sampling seasons in Fehérvárcsurgó reservoir, while the amplicons of the sxtS gene were not observed in our study. Despite the absence of the sxtS gene, the presence of previously reported STX producers A. flos-aque (Ferreira et al., 2001) and C. issatschenkoi (Li et al., 2003; Pereira et al., 2000) in the reservoir raises concerns over the potential production of STXs. The sxt gene cluster is genus-specific and comprises about 30 genes (Mihali et al., 2009),



554 Page 10 of 14 Environ Monit Assess (2021) 193:554



**Fig. 5** Histopathological sections of intestine (**A**, **B**), spleen (**C**, **D**), testis (**E**, **F**), ovaries (**G**, **H**), and muscles (**I**, **J**) of controls (**A**, **C**, **E**, **G**, **I**) and individuals sampled from the Fehé-

rvárcsurgó reservoir during 2018 (**B**, **D**, **F**, **H**, **J**). H&E staining. **G**, **H** 200 μm; **A**, **B**, **E**, **F** 100 μm; **C**, **D**, **I**, **J** 50 μm

four of which (*sxtA*, *sxtB*, *sxtG*, *sxtS*) are conserved among all STX-producing genera. The presence of all genes within the cluster is necessary for the synthesis of STXs. The observed lack of amplification of the *sxtS* gene may be due to mutations within the gene cluster leading to non-functional gene clusters and an inability of the observed strains to produce STXs.

Several fish tissues were also tested for the presence of MCs but accumulation was not observed. Cyanotoxins could have remained covalently bound to protein phosphatases in the tissue (Ibelings et al., 2005; MacKintosh et al., 1990; Williams et al., 1997a, b), excreted and concentrations reduced by detoxification (Adamovský et al., 2007; Malbrouck & Kestemont, 2006; Snyder et al., 2002; Xie et al., 2004), or concentrations in water were not sufficient enough to be accumulated in detectable levels. Similar conclusions were reached during the research conducted on Lake Ludoš in the neighboring country Serbia during 2018, when cyanotoxins in fish tissue were not detected (Tokodi et al., 2020). However, at the same site a few years earlier (2011/2012), the accumulation of MC-RR and MC-LR in the tissue samples of the Prussian carp from Lake Ludoš was documented, alongside with histopathological changes in liver, kidney, gills, and intestine of exposed fish (Tokodi et al., 2018). Another study from fishponds in Serbia has also reported MC-RR accumulation in muscle tissues and histopathological changes in muscles, gills, intestines, kidneys, and liver of *Cyprinus carpio* farmed in blooming waters (Drobac et al., 2016). Previous studies used a different analytical method (Drobac et al., 2016; Tokodi et al., 2018) better suited for testing accumulation of cyanotoxins in fish samples, which could potentially explain lack of data; however, the method that was used during this investigation was previously successfully used with spiked samples before fish tissue preparation which resulted in positive detection of investigated MCs (Tokodi et al., 2020).

Histopathological changes were also noted in fish caught from Fehérvárcsurgó reservoir. Alterations were more severe during May and especially July, which coincides with the increase in cyanobacterial biomass and the detection of MC-RR in July. However, the absence of hepatic damage indicates that MCs were not the main or only damage-causing agents. Noted vacuolization often occurs due to exposure of fish and mammals to cyanotoxins (Carbis et al., 1996; Drobac et al., 2016; Fischer & Dietrich, 2000; Gupta et al., 2003; Hooser et al., 1990; Jiang et al., 2011; Li et al., 2007; Svirčev et al., 2015); however, this change is very nonspecific and can occur due to the action of various agents, but also due to stress, or dietary imbalance (Thoolen et al., 2010). Also, the alterations in liver observed in April may be the result of cyanobacterial blooms that occurred



prior to this study, and it is possible that these changes are accumulations of previous toxin action. The changes observed in the kidneys were more intense than the changes in the liver. Given that Fischer and Dietrich (2000) observed that cyprinid fish display more intense changes in the kidneys rather than the liver after exposure to MCs, and the fact that the kidneys have membrane transporters for MC, it is possible that alterations were caused by MCs. However, since very low concentrations of MCs were detected, it is not clear whether these concentrations are sufficient to cause kidney damage. Most of the mentioned alterations in gills represent defense mechanisms which reduce the respiratory surface and thus reduce the uptake/diffusion of toxic compounds (Lujić et al., 2015). These changes occur after exposure of fish to cyanotoxins (Svirčev et al., 2015) and are most often the result of acute toxin exposure. However, due to the complex relationships that occur within the ecosystem, it is possible that the observed changes are the result of chronic exposure to certain toxins that were not measured in this study but were synthesized by cyanobacteria present in the ecosystem. Since the changes in the liver, kidneys, and gills of fish from this Fehérvárcsurgó reservoir were very mild compared to some other ecosystems (Marinović et al., 2021; Tokodi et al., 2020), and that the cyanobacterial blooming was less intense, the absence of histopathological alterations in the intestines, spleen, testicles, ovaries, and muscles was expected.

Humans can come into contact with cyanobacteria and their metabolites through, e.g., skin and mucosal tissue contact, ingestion of water and food, or aerosol inhalation. Given the multipurpose use of the Fehérvárcsurgó reservoir, fishing, swimming, recreation, and agricultural practices all lead to exposure scenarios. An example from China corroborates the potential threats of MCs in lake/reservoir water: fishermen with chronic and combined exposure to cyanobacteria (contaminated water, fish, and other aquatic organisms from blooming lakes) had serious health problems in the form of hepatocellular damage, and MCs were identified in their serum (Chen et al., 2009). Previously published data indicate the toxic blooming of cyanobacteria in the Fehérvárcsurgó reservoir occurred in the past; however, so far, there is no publication that clearly describes the blooming, production of toxins, or the consequences it has to fish population. Recently, similar investigation was performed in Hungary on Lake Balaton and KBWPS (Marinović et al., 2021). During 2018, blooming of toxic and potentially toxic cyanobacterial species was noted, as well as five MC congeners. Histopathological alterations of the fish from KBWPS were more severe than those described in this study, which could be explained by higher severity of the blooming. Results also showed potential blooming problem throughout the year, as it was the case in this study, which demonstrates necessity of regular monitoring of the cyanobacterial and cyanotoxin distribution, so that potential problems can be anticipated and prevented before escalation.

#### Conclusions

Research performed on the Fehérvárcsurgó reservoir in Hungary during 2018 showed that cyanobacteria were present and blooming during sampling in April, May, July, and September. The highest number of cells per mL was recorded in July (2,822,000), and the dominant species was Aphanizomenon flos-aquae (2,124,000 cells/mL in July). Microcystis flos-aquae, Microcystis wesenbergii, Cuspidothrix issatschenkoi, Dolichospermum flos-aquae, and Snowella litoralis were also detected, and their abundance varied depending on the season. Analyses of cyanotoxin encoding genes showed amplification of MC (mcyE) and STX (sxtG) coding genes, while cyanotoxin analyses indicated a low concentration of MC-RR in July. The identification of MC-RR and cyanotoxin encoding genes indicated that MCs and STXs pose a latent threat that could cause future problems for the reservoir ecosystem and its users, such as fishermen and swimmers. Finally, the analyses of fish tissue did not show the presence of the investigated MCs, but several histopathological alterations, predominantly in gills and kidneys, were observed in fish caught from the reservoir. This type of research can uncover hidden dangers posed by cyanobacteria, and thus give us the advantage in preventing potential consequences and preserving the balance of the ecosystem.

Acknowledgements The authors would like to thank Henna Savela for the supervision of molecular analysis. The equipment and facilities for all molecular biology work were provided by the Department of Life Technologies/Biotechnology, University of Turku. The authors would like to express gratitude to Árpád Ferincz and Ádám Staszny for assistance in field sampling.



Author contribution DDB conceived and designed the study, collected the samples, performed the detection of cyanotoxins, and wrote the manuscript. NT conceived and designed the study, collected the samples, performed the detection of cyanotoxins, material preparation, data collection, analysis and interpretation, and functions as corresponding author. TD performed the PCR analysis of cyanotoxin encoding genes. SS and NĐ performed the qualitative and quantitative analyses of cyanobacteria. ZM, JL, NK, and IŠ collected the fish samples and analyzed the histology of fish tissues. JL conceived the study and with BU supervised the project. JM and ZS were study advisors and revised the manuscript critically. All the authors read and approved the final manuscript.

**Funding** The authors received financial support from the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant No. 451–03-9/2021–14/200125 and No. 451–03-9/2021–14/200122), Bilateral project Hungary-Serbia Invasive and blooming cyanobacteria in Serbian and Hungarian waters (2017–2019) (TET\_16-1–2016-0176) and the Erasmus+programme of the European Union (agreement number: 2017–1-FI01-KA107-034440). The Åbo Akademi University Doctoral Scholarship Program supported the work performed by TD.

**Availability of data and material** The data that support the findings of this study are available from the corresponding author, NT, upon reasonable request.

## **Declarations**

**Ethics approval** Collecting adult fish from a natural population was approved by the Hungarian Ministry of Agriculture (license HaGF/113/2019).

**Competing interests** The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

# References

Adamovský, O., Kopp, R., Hilscherova, K., Babica, P., Palikova, M., Paskova, V., Navratil, S., & Blaha, L. (2007). Microcystin

- kinetics (bioaccumulation, elimination) and biochemical responses in common carp and silver carp exposed to toxic cyanobacterial blooms. *Environmental Toxicology and Chemistry*, 26, 2687–2693.
- Ballot, A., Scherer, P. I., & Wood, S. A. (2018). Variability in the anatoxin gene clusters of *Cuspidothrix issatschenkoi* from Germany, New Zealand, China and Japan. *PLoS One*, 13(7), e0200774.
- Bernard, C., Ballot, A., Thomazeau, S., Maloufi, S., Furey, A., Mankiewicz-Boczek, J., Pawlik-Skowrońska, B., Capelli, C., & Salmaso, N. (2017). Appendix 2: cyanobacteria associated with the production of cyanotoxins. In J. Meriluoto, L. Spoof., G.A. Codd (Eds.) *Handbook of cyanobacterial* monitoring and cyanotoxin analysis (pp. 501–525). John Wiley & Sons, Ltd.
- Bouaïcha, N., Miles, C., Beach, D., Labidi, Z., Djabri, A., Benayache, N., & Nguyen-Quang, T. (2019). Structural diversity, characterization and toxicology of microcystins. *Toxins (Basel)*, 11, 714. https://doi.org/10.3390/ toxins11120714
- Carbis, C. R., Rawlin, G. T., Mitchell, G. F., Anderson, J. W., & McCauley, I. (1996). The histopathology of carp, *Cyprinus carpio* L., exposed to microcystins by gavage, immersion and intraperitoneal administration. *Journal of Fish Disease*, 19, 199–207. https://doi.org/10.1111/j. 1365-2761.1996.tb00126.x
- Chen, J., Xie, P., Li, L., & Xu, J. (2009). First identification of the hepatotoxic microcystins in the serum of a chronically exposed human population together with indication of hepatocellular damage. *Toxicological Sciences*, 108, 81–89.
- Dittman, E., Mankiewicz-Boczek, J., & Gagała, I. (2017). SOP 6.2: PCR detection of microcystin and nodularin biosynthesis genes in the cyanobacterial orders Oscillatoriales, Chroococcales, Stigonematales, and Nostocales. In R. Kurmayer, K. Sivonen, A.Wilmotte, N. Salmaso (Eds.), Molecular Tools for the Detection and Quantification of Toxigenic Cyanobacteria (pp. 175–178). John Willey & Sons, Ltd.: Chichester, UK.
- Drobac, D., Tokodi, N., Lujić, J., Marinović, Z., Subakov-Simić, G., Dulić, T., Važić, T., Nybom, S., Meriluoto, J., Codd, G. A., & Svirčev, Z. (2016). Cyanobacteria and cyanotoxins in fishponds and their effects on fish tissue. *Harmful Algae*, 55, 66–76.
- Falconer, I. R. (1998). Algal Toxins and Human Health. In J. Hrubec (Ed.), *The Handbook of Environmental Chemistry 5. Part C Quality and Treatment of Drinking Water II*. (pp. 53–82). Berlin, Heidelberg: Springer-Verlag.
- Ferreira, F. M. B., Soler, J. M. F., Fidalgo, M. L., & Fernández-Vila, P. (2001). PSP toxins from *Aphanizomenon flos-aquae* (cyanobacteria) collected in the Crestuma-Lever reservoir (Douro river, northern Portugal). *Toxicon*, 39, 757–761.
- Fischer, W. J., & Dietrich, D. R. (2000). Pathological and biochemical characterization of microcystin-induced hepatopancreas and kidney damage in carp (*Cyprinus carpio*). *Toxicology and Applied Pharmacology*, 164, 73–81. https://doi.org/10.1006/taap.1999.8861
- Garcia-Pichel, F., Belnap, J., Neuer, S., & Schanz, F. (2003). Estimates of global cyanobacterial biomass and its distribution. *Algological Studies*, 109(1), 213–227.



Environ Monit Assess (2021) 193:554 Page 13 of 14 554

Gupta, N., Pant, S. C., Vijayaraghavan, R., & Rao, P. V. L. (2003). Comparative toxicity evaluation of cyanobacterial cyclic peptide toxin microcystin variants (LR, RR, YR) in mice. *Toxicology*, 188, 285–296. https://doi.org/10.1016/ S0300-483X(03)00112-4

- Hodoki, Y., Ohbayashi, K., Kobayashi, Y., Okuda, N., & Nakano, S. (2012). Detection and identification of potentially toxic cyanobacteria: Ubiquitous distribution of *Microcystis aeruginosa* and *Cuspidothrix issatschenkoi* in Japanese lakes. *Harmful Algae*, 16, 49–57.
- Hooser, S. B., Beasley, V. R., Basgall, E. J., Carmichael, W. W., & Haschek, W. M. (1990). Microcystin-LR-induced ultrastructural changes in rats. *Veterinary Pathology*, 27, 9–15. https://doi.org/10.1177/030098589002700102
- Ibelings, B. W., Bruning, K., de Jonge, J., Wolfstein, K., & Pires, L. D. M. (2005). Distribution of microcystins in a lake food web: No evidence for biomagnification. *Microbial Ecology*, 49, 487–500.
- Jasprica, N., Hafner, D., Batistić, M., & Kapetanović, T. (2005). Phytoplankton in three freshwater lakes in the Neretva River delta (Eastern Adriatic, NE Mediterranean). Nova Hedwigia, 81, 37–54.
- Jiang, J., Gu, X., Song, R., Zhang, Q., Geng, J., Wang, X., & Yang, L. (2011). Time-dependent oxidative stress and histopathological changes in *Cyprinus carpio* L. exposed to microcystin-LR. *Ecotoxicology*, 20, 1000–1009. https://doi.org/10.1007/s10646-011-0646-9
- Komárek, J. (2013). Cyanoprokaryota 3. Teil: Heterocytous Genera. In B. Budel, G. Gärtner, L. Krienitz, M. Schagerl (Eds.), Süβwasserflora von Mitteleuropa. (pp. 1–1130). Springer Spektrum Verlag: Heidelberg/Berlin, Germany.
- Komárek, J., & Anagnostidis, K. (1998). Cyanoprokaryota 1. Teil: Chroococcaless. In H. Ettl, J. Gerloff, H. Heynig, D. Mollenhauer, (Eds.), Süβwasserflora von Mitteleuropa. (pp. 1–548). Spektrum Akademischer Verlag: Heidelberg/Berlin,Germany.
- Komárek, J., & Anagnostidis, K. (2005). Cyanoprokaryota 2. Teil: Oscillatoriales. In B. Budel, L. Krienitz, G. Gärtner, M. Schagerl (Eds.), Süβwasserflora von Mitteleuropa. (pp. 1–759). Spektrum Akademischer Verlag: Heidelberg/Berlin, Germany.
- Li, R., Carmichael, W. W., & Pereira, P. (2003). Morphological and 16s rRNA gene evidence for reclassification of the paralytic shellfish toxin producing *Aphanizomenon flos-aquae Imecya* 31 as *Aphanizomenon issatschenkoi* (cyanophyceae). *Journal of Phycology*, 39, 814–818.
- Li, L., Xie, P., & Chen, J. (2007). Biochemical and ultrastructural changes of the liver and kidney of the phytoplank-tivorous silver carp feeding naturally on toxic *Microcystis* blooms in Taihu Lake, China. *Toxicon*, 49, 1042–1053. https://doi.org/10.1016/j.toxicon.2007.01.013
- Lujić, J., Matavulj, M., Poleksić, V., Rašković, B., Marinović, Z., Kostić, D., & Miljanović, B. (2015). Gill reaction to pollutants from the Tamiš River in three freshwater fish species, Esox lucius L. 1758, Sander lucioperca (L. 1758) and Silurus glanis L. 1758: A comparative study. Anatomia Histologia Embryologia, 44, 128–137. https://doi.org/10.1111/ahe.12119
- Lyon-Colbert, A., Su, S., & Cude, C. (2018). A systematic literature review for evidence of *Aphanizomenon flos-aquae*

- toxigenicity in recreational waters and toxicity of dietary supplements: 2000–2017. *Toxins (Basel)*, 10(7), 254.
- MacKintosh, C., Beattie, K. A., Klumpp, S., Cohen, P., & Codd, G. A. (1990). Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from both mammals and higher plants. *FEBS Letters*, 264, 187–192.
- Malbrouck, C., & Kestemont, P. (2006). Effects of microcystins on fish. Environmental Toxicology and Chemistry, 25, 72–86
- Marinović, Z., Tokodi, N., Backović, D. D., Šćekić, I., Kitanović, N., Simić, S. B., Đorđević, N. B., Ferincz, Á., Staszny, Á., Dulić, T., Meriluoto, J., Urbányi, B., Lujić, J., & Svirčev, Z. (2021). Does the Kis-Balaton Water Protection System (KBWPS) effectively safe-guard Lake Balaton from toxic cyanobacterial blooms? *Microorganisms*, 9, 960. https://doi.org/10.3390/microorganisms9050960
- Mazmouz, R., Chapuis-Hugon, F., Mann, S., Pichon, V., Méjean, A., & Ploux, O. (2010). Biosynthesis of cylindrospermopsin and 7-epicylindrospermopsin in *Oscillatoria* sp. strain PCC 6506: Identification of the *cyr* gene cluster and toxin analysis. *Applied Environmental Microbiology*, 76, 4943–4949.
- Mihali, T. K., Kellmann, R., & Neilan, B. A. (2009). Characterisation of the paralytic shellfish toxin biosynthesis gene clusters in *Anabaena circinalis* AWQC131C and *Aphanizomenon* sp. NH 5. *BMC Biochemistry*, 10, 8.
- Nagy, S., Csőke, B., Zajzon, N., Kristály, F., Márkus, I. R., Gombkoto, I., Pap, Z., Kaliczné Papp, K., Szép, L., Duber, N., Schupler, H., & Gaga, I. (2014). Heavy minerals from glass sand tailings, Fehérvárcsurgó. In J. Földessy (Ed.), Basic research of the strategic raw materials in Hungary (pp. 120–150). Millagrossa Kft.
- Otsuka, S., Suda, S., Li, R., Watanabe, M., Oyaizu, H., Matsumoto, S., & Watanabe, M. M. (1999). Phylogenetic relationships between toxic and non-toxic strains of the genus *Microcystis* based on 16S to 23S internal transcribed spacer sequence. *FEMS Microbiology Letters*, 172(1), 15–21.
- Pereira, P., Onodera, H., Andrinolo, D., Franca, S., Araújo, F., Lagos, N., & Oshima, Y. (2000). Paralytic shellfish toxins in the freshwater cyanobacterium *Aphanizomenon flos-aquae*, isolated from Montargil reservoir, Portugal. *Toxicon*, 38, 1689–1702.
- Rajaniemi-Wacklin, P., Rantala, A., Mugnai, M. A., Turicchia, S., Ventura, S., Komárková, J., Lepistö, L., & Sivonen, K. (2006). Correspondence between phylogeny and morphology of Snowella spp. and Woronichinia naegeliana, cyanobacteria commonly occurring in lakes. *Journal of Phycology*, 42(1), 226–232.
- Rantala-Ylinen, A., Känä, S., Wang, H., Rouhiainen, L., Wahlsten, M., Rizzi, E., Berg, K., Gugger, M., & Sivonen, K. (2011). Anatoxin-a synthetase gene cluster of the cyanobacterium Anabaena sp. strain 37 and molecular methods to detect potential producers. Journal of Applied Environmental Microbiology, 77, 7271–7278.
- Savela, H., Spoof, L., Perälä, N., Preede, M., Lamminmäki, U., Nybom, S., Häggqvist, K., Meriluoto, J., & Vehniäinen, M. (2015). Detection of cyanobacteria sxt genes and paralytic shellfish toxins in freshwater lakes and brakish waters on Åland Islands, Finland. *Harmful Algae*, 46, 1–10.
- Savela, H., Spoof, L., Höysniemi, N., Vehniäinen, M., Mankiewicz-Boczek, J., Jurczak, T., Kokociński, M., &



554 Page 14 of 14 Environ Monit Assess (2021) 193:554

Meriluoto, J. (2017). First report of cyanobacterial paralytic shellfish toxin biosynthesis genes and paralytic shellfish toxin production in Polish freshwater lakes. *Advances in Oceanography and Limnology*, 8, 61–70.

- Snyder, G. S., Goodwin, A. E., & Freeman, D. W. (2002). Evidence that channel catfish, *Ictalurus punctatus* (Rafinesque), mortality is not linked to ingestion of the hepatotoxin microcystin-LR. *Journal of Fish Diseases*, 25, 275–286.
- Stoyneva-Gärtner, M., Stefanova, K., Descy, J.-P., Uzunov, B., Radkova, M., Pavlova, V., Mitreva, M., & Gärtner, G. (2021). *Microcystis aeruginosa* and *M. wesenbergii* were the primary planktonic microcystin producers in several Bulgarian waterbodies (August 2019). *Applied Sciences*, 11, 357. https://doi.org/10.3390/app11010357
- Svirčev, Z., Lujić, J., Marinović, Z., Drobac, D., Tokodi, N., Stojiljković, B., & Meriluoto, J. (2015). Toxicopathology induced by microcystins and nodularin: A histopathological review. *Journal of Environmental Science and Health, Part C*, 33, 125–167.
- Svirčev, Z., Drobac, D., Tokodi, N., Mijović, B., Codd, G. A., & Meriluoto, J. (2017). Toxicology of microcystins with reference to cases of human intoxications and epidemiological investigations of exposures to cyanobacteria and cyanotoxins. Archives of Toxicology, 91(2), 621–650.
- Svirčev, Z., Lalić, D., Bojadžija Savić, G., Tokodi, N., Drobac Backović, D., Chen, L., Meriluoto, J., & Codd, G. A. (2019). Global geographical and historical overview of cyanotoxin distribution and cyanobacterial poisonings. Archives of Toxicology, 93, 2429–2481.
- Thoolen, B., Maronpot, R. R., Harada, T., Nyska, A., Rousseaux, C., Nolte, T., Malarkey, D. E., Kaufmann, W., Küttler, K., Deschl, U., Nakae, D., Gregson, R., Vinlove, M. P., Brix, A. E., Singh, B., Belpoggi, F., & Ward, J. M. (2010). Proliferative and nonproliferative lesions of the rat and mouse hepatobiliary system. *Toxicologic Pathology*, 38, 5–81. https://doi.org/10.1177/0192623310386499
- Tokodi, N., Drobac, D., Meriluoto, J., Lujić, J., Marinović, Z., Važić, T., Nybom, S., Simeunović, J., Dulić, T., Lazić, G., Petrović, T., Vuković-Gačić, B., Sunjog, K., Kolarević, S., Kračun-Kolarević, M., Subakov-Simić, G., Miljanović, B., Codd, G. A., & Svirčev, Z. (2018). Cyanobacterial effects in Lake Ludoš, Serbia is preservation of a degraded aquatic ecosystem justified? Science of Total Environment, 635, 1047–1062.

- Tokodi, N., Drobac Backović, D., Lujić, J., Šćekić, I., Simić, S., Đorđević, N., Dulić, T., Miljanović, B., Kitanović, N., Marinović, Z., Savela, H., Meriluoto, J., & Svirčev, Z. (2020). Protected freshwater ecosystem with incessant cyanobacterial blooming awaiting a resolution. Water, 12, 129. https://doi.org/10.3390/w12010129
- Törökné, A. (1999). A new culture-free microbiotest for routine detection of cyanobacterial toxins. *Environmental Toxicology*, *14*(5), 466–472.
- Törökné, A., Palovics, A., & Bankine, M. (2001). Allergenic (sensitization, skin and eye irritation) effects of freshwater cyanobacteria-experimental evidence. *Environmental Toxicology*, 16, 512–516.
- Utermöhl, H. (1958). Zurvervolkmmung der quantitativen phytoplankton methodik. *Mitteilungen Internationale Vereinigung Fur Theoretische Und Angewandte Limnologie*, *9*, 1–38.
- Via- Ordorika, L., Fastner, J., Kurmayer, R., Hisbergues, M., Dittmann, E., Komarek, J., Erhard, M., & Chorus, I. (2004). Distribution of microcystin-producing and non-microcystin-producing *Microcystis* sp. in European Freshwater bodies: Detection of microcystins and microcystin genes in individual colonies system. *Applied Microbiology*, 27, 592–602.
- Williams, D. E., Craig, M., Dawe, S. C., Kent, M. L., Andersen, R. J., & Holmes, C. F. B. (1997a). C-14-Labeled microcystin-LR administered to *Atlantic salmon* via intraperitoneal injection provides in vivo evidence for covalent binding of microcystin-LR in salmon livers. *Toxicon*, 35, 985–989.
- Williams, D. E., Craig, M., Dawe, S. C., Kent, M. L., Holmes, C. F. B., & Andersen, R. J. (1997b). Evidence for a covalently bound form of microcystin-LR in salmon liver and Dungeness crab larvae. *Chemical Research in Toxicology*, 10, 463–469.
- Xie, L., Xie, P., Ozawa, K., Honma, T., Yokoyama, A., & Park, H. D. (2004). Dynamics of microcystins-LR and -RR in the phytoplanktivorous silver carp in a sub-chronic toxicity experiment. *Environmental Pollution*, 127, 431–439.
- Xu, Y., Wu, Z., Yu, B., Peng, X., Yu, G., Wei, Z., Wang, G., & Li, R. (2008). Non-microcystin producing *Microcystis* wesenbergii (Komárek) Komárek (Cyanobacteria) representing a main waterbloom - forming species in Chinese waters. Environmental Pollution, 156(1), 162–167.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

