Energy of Graphs with Self-Loops

Ivan Gutman^{1,*}, Izudin Redžepović¹, Boris Furtula¹, Ali Mohammed Sahal²

¹Faculty of Science, University of Kragujevac, P.O.Box 60, 34000 Kragujevac, Serbia

 $gutman@kg.ac.rs \ , \ izudin.redzepovic@pmf.kg.ac.rs \ , \ furtula@uni.kg.ac.rs$

²Department of Studies in Mathematics, Faculty of Education – Zabid, University of Hodeidah, Republic of Yemen alisahl1980@gmail.com

(Received August 2, 2021)

Abstract

The energy of graphs containing self-loops is considered. If the graph G of order n contains σ self-loops, then its energy is defined as $E(G) = \sum |\lambda_i - \sigma/n|$ where $\lambda_1, \lambda_2, \ldots, \lambda_n$ are the eigenvalues of the adjacency matrix of G. Some basic properties of E(G) are established, and several open problems pointed out or conjectured.

1 Introduction

A graph is said to be simple (or schlicht) if it does not possess directed, weighted or multiple edges, and self-loops [12,13,21]. Let G be a simple graph of order n, with vertex set $\mathbf{V} = \{v_1, v_2, \dots, v_n\}$. Its adjacency matrix $\mathbf{A}(G)$ is a square symmetric matrix of order n whose (i, j)-element is defined as

$$\mathbf{A}(G)_{ij} = \begin{cases} 1 & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent,} \\ 0 & \text{if the vertices } v_i \text{ and } v_j \text{ are not adjacent,} \\ 0 & \text{if } i = j. \end{cases}$$

Let $\lambda_1(G), \lambda_2(G), \ldots, \lambda_n(G)$ be the eigenvalues of $\mathbf{A}(G)$. Then the energy of G is

$$E(G) = \sum_{i=1}^{n} |\lambda_i(G)|.$$
 (1)

^{*}Corresponding author

The theory of graph energy is nowadays a well elaborated field of applied mathematics and mathematical chemistry [14,18]. One should recall that the concept of graph energy has a chemical origin and a chemical interpretation [8].

There are more than a thousand papers on graph energy and its variants [9, 10]. Practically all these papers are concerned with simple graphs. It is remarkable that the energy of graph with self-loops has not been considered so far. This is additionally surprising because graphs with self-loops (representing heteroatoms) are of evident chemical significance, and were much studied in the past, including their spectral properties [1, 5, 6, 15, 16, 19].

Let **S** be a subset of V(G). The number of elements of **S** will be denoted by σ .

Let G_S be the graph obtained from the simple graph G, by attaching a self-loop to each of its vertices belonging to **S**. Then the adjacency matrix of G_S is a symmetric square matrix $\mathbf{A}(G_S)$ of order n, whose (i, j)-element is defined as

$$\mathbf{A}(G_S)_{ij} = \begin{cases} 1 & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent,} \\ 0 & \text{if the vertices } v_i \text{ and } v_j \text{ are not adjacent,} \\ 1 & \text{if } i = j \text{ and } v_i \in \mathbf{S}, \\ 0 & \text{if } i = j \text{ and } v_i \notin \mathbf{S}. \end{cases}$$

Let $\lambda_1(G_S), \lambda_2(G_S), \ldots, \lambda_n(G_S)$ be the eigenvalues of $\mathbf{A}(G_S)$. These all are real-valued, and

$$\sum_{i=1}^{n} \lambda_i(G_S) = \sigma \,. \tag{2}$$

Therefore, the energy of G_S (in full analogy to the energy of any matrix with non-zero diagonal [2,3,11]) has to be defined as

$$E(G_S) = \sum_{i=1}^{n} \left| \lambda_i(G_S) - \frac{\sigma}{n} \right| \tag{3}$$

In this paper, we begin to examine the properties of $E(G_S)$.

2 Basic properties of energy of graphs with self-loops

First we state an elementary result:

Lemma 1. (a) If $\sigma = 0$, then $E(G_S) = E(G)$. (b) If $\sigma = n$, then $E(G_S) = E(G)$. *Proof.* Lemma 1(a) is trivially obvious, since for $\sigma = 0$, the graphs G_S and G coincide.

Let \mathbf{I}_n denote the unit matrix of order n. If $\sigma = n$, then $\mathbf{A}(G_S) = \mathbf{A}(G) + \mathbf{I}_n$. Therefore, $\lambda_i(G_S) = \lambda_i(G) + 1$, i = 1, 2, ..., n. Lemma 1(b) follows then from Eqs. (1) and (3).

Lemma 1 immediately leads to the question: What is the relation between $E(G_S)$ and E(G) in the case of $1 \le \sigma \le n - 1$? Based on our numerical studies, we state the following claim:

Conjecture 2. Let G be any simple graph of order n, and **S** any subset of its vertices, $1 \le \sigma \le n - 1$. Then $E(G_S) > E(G)$.

If this conjecture is true, then we are faced with many further questions. For instance, for a given graph (or class of graphs), for which **S** is the difference $E(G_S) - E(G)$ maximal? Etc. etc.

Some simple examples, illustrating Conjecture 2 are collected in Table 1.

S	$E(G_S)$
Ø	4.4721
$\{v_1\}$	4.7588
$\{v_2\}$	4.6659
$\{v_1, v_2\}$	4.6056
$\{v_1, v_3\}$	4.9770
$\{v_1, v_4\}$	4.8284
$\{v_2, v_3\}$	4.8284
$\{v_1, v_2, v_3\}$	4.7588
$\{v_1, v_2, v_4\}$	4.6659
$\{v_1, v_2, v_3, v_4\}$	4.4721

Table 1. Energies of the path P_4 with self-loops.

From Table 1 we see that the energy depends not only on the number of self-loops, but also on their position. We, however, observe that the energies of some different graphs coincide. This is the consequence of the following general result.

Theorem 3. Let G be a bipartite graph of order n, with vertex set V. Let S be a subset of V. Then $E(G_S) = E(G_{V \setminus S})$.

Proof. We prove Theorem 3 assuming that n is even. If n is odd, the proof would proceed in a fully analogous manner.

For the proof of Theorem 3, we need to recall details of the Sachs coefficient theorem [1, 4, 7, 20].

Let $\phi(G_S, \lambda) = \det [\lambda \mathbf{I}_n - \mathbf{A}(G_S)]$ be the characteristic polynomial of G_S . We write it in the form

$$\phi(G_S, \lambda) = \phi_e(G_S, \lambda) + \phi_o(G_S, \lambda) \tag{4}$$

where

$$\phi_e(G_S, \lambda) = \sum_{k \ge 0} c_{2k} \,\lambda^{n-2k} \qquad \text{and} \qquad \phi_o(G_S, \lambda) = \sum_{k \ge 0} c_{2k+1} \,\lambda^{n-2k-1}$$

Let \mathbf{H}_k be the set of k-vertex subgraphs of G_S whose components are cycles and/or copies of P_2 and/or isolated vertices with self-loops. According to the Sachs theorem,

$$c_k = \sum_{H \in \mathbf{H}_k} (-1)^{p(H)} 2^{c(H)}$$
(5)

where p(H) is the number of components of H and c(H) the number of cycles of H.

Since G_S is bipartite, all its cycles (if any) are of even size. Therefore, all subgraphs contained in H_{2k} must possess and even number (or zero) of vertices with self-loops. All components of H_{2k+1} must possess an odd number of vertices with self-loops.

Let G_S^- be the graph obtained from G_S by changing the signs of all its self-loops from +1 to -1. Then by the Sachs formula (5),

$$\phi(G_S^-, \lambda) = \phi_e(G_S, \lambda) - \phi_o(G_S, \lambda).$$

Let ξ be a zero of the polynomial $\phi(G_S, \lambda)$. Then in view of Eq. (4),

$$\phi_e(G_S,\xi) = -\phi_o(G_S,\xi) \,.$$

In addition,

$$\phi_e(G_S,\xi) = \phi_e(G_S,-\xi) \quad \text{and} \quad -\phi_o(G_S,\xi) = \phi_o(G_S,-\xi)$$

implying

$$\phi_e(G_S, -\xi) - \phi_o(G_S, -\xi) = 0$$
 i.e., $\phi(G_S^-, -\xi) = 0$

We thus conclude that if $\lambda_1(G_S), \lambda_2(G_S), \ldots, \lambda_n(G_S)$ are the eigenvalues of G_S , then $-\lambda_1(G_S), -\lambda_2(G_S), \ldots, -\lambda_n(G_S)$ are the eigenvalues of G_S^- .

Consider now the characteristic polynomial of $G_{V \setminus S}$.

Let $\mathbf{A}(G_S) = \mathbf{A}(G) + \mathbf{J}_S$ and $\mathbf{A}(G_{V \setminus S}) = \mathbf{A}(G) + \mathbf{J}_{V \setminus S}$, with $\mathbf{J}_S + \mathbf{J}_{V \setminus S} = \mathbf{I}_n$. Then

$$\phi(G_{V\setminus S}, \lambda) = \det \left[\lambda \mathbf{I}_n - \mathbf{A}(G) - \mathbf{J}_{V\setminus S}\right] = \det \left[\lambda \mathbf{I}_n - \mathbf{A}(G) - \mathbf{I}_n + \mathbf{J}_S\right]$$
$$= \det \left[(\lambda - 1) \mathbf{I}_n - (\mathbf{A}(G) - \mathbf{J}_S)\right] = \det \left[(\lambda - 1) \mathbf{I}_n - \mathbf{A}(G_S^-)\right]$$

from which it follows

$$\phi(G_{V\setminus S},\lambda) = \phi(G_S^-,\lambda-1).$$

This means that the eigenvalues $\lambda_i(G_{V\setminus S})$, i = 1, 2, ..., n, coincide with $\lambda_i(G_S^-) + 1$, i = 1, 2, ..., n. Bearing this in mind, we have

$$E(G_{V\setminus S}) = \sum_{i=1}^{n} \left| \lambda_i(G_{V\setminus S}) - \frac{n-\sigma}{n} \right| = \sum_{i=1}^{n} \left| \lambda_i(G_{V\setminus S}) - 1 + \frac{\sigma}{n} \right|$$
$$= \sum_{i=1}^{n} \left| \lambda_i(G_S^-) + 1 - 1 + \frac{\sigma}{n} \right| = \sum_{i=1}^{n} \left| -\lambda_i(G_S) + \frac{\sigma}{n} \right| = \sum_{i=1}^{n} \left| \lambda_i(G_S) - \frac{\sigma}{n} \right|.$$

Theorem 3 follows now by Eq. (3).

The equality $E(G_S) = E(G_{V\setminus S})$ does not hold if the graph G is not bipartite. The simplest example showing this is the triangle with one self-loop (whose energy is 4.1618) and with two self-loops (whose energy is 4.1308).

3 McClelland-type bound for the energy of graphs with self-loops

In this section we obtain a McClelland–type upper bound [17] for the energy of graphs with self-loops. In order to achieve this goal, we first establish a few auxiliary results.

Lemma 4. Let G_S be a graph of order n, with m edges, and σ self-loops. Let $\lambda_1, \lambda_2, \ldots, \lambda_n$ be its eigenvalues. Then

$$\sum_{i=1}^n \lambda_i^2 = 2m + \sigma \,.$$

Proof.

$$\sum_{i=1}^{n} \lambda_{i}^{2} = \sum_{i=1}^{n} \left[\mathbf{A}(G_{S})^{2} \right]_{ii} = \sum_{i=1}^{n} \left[\left(\mathbf{A}(G) + \mathbf{J}_{S} \right)^{2} \right]_{ii}$$
$$= \sum_{i=1}^{n} \left[\mathbf{A}(G)^{2} + \mathbf{A}(G) \mathbf{J}_{S} + \mathbf{J}_{S} \mathbf{A}(G) + \mathbf{J}_{S}^{2} \right]_{ii}.$$

By direct calculation it is easy to shown that

$$\sum_{i=1}^{n} \left[\mathbf{A}(G)^{2} \right]_{ii} = 2m \ , \ \sum_{i=1}^{n} \left[\mathbf{A}(G) \, \mathbf{J}_{S} \right]_{ii} = \sum_{i=1}^{n} \left[\mathbf{J}_{S} \, \mathbf{A}(G) \right]_{ii} = 0 \ , \ \sum_{i=1}^{n} \left[\mathbf{J}_{S}^{2} \right]_{ii} = \sigma$$

from which Lemma 4 follows.

Lemma 5. With the same notation as in Lemma 4,

$$\sum_{i=1}^{n} \left| \lambda_i - \frac{\sigma}{n} \right|^2 = 2m + \sigma - \frac{\sigma^2}{n}$$

Proof.

$$\sum_{i=1}^{n} \left| \lambda_i - \frac{\sigma}{n} \right|^2 = \sum_{i=1}^{n} \left(\lambda_i^2 - 2\lambda_i \frac{\sigma}{n} + \frac{\sigma^2}{n^2} \right) = \sum_{i=1}^{n} \lambda_i^2 - 2\frac{\sigma}{n} \sum_{i=1}^{n} \lambda_i + \frac{\sigma^2}{n^2}$$

and Lemma 5 follows by using Lemma 4 and formula (2).

Theorem 6. Let G_S be a graph of order n, with m edges, and σ self-loops. Then

$$E(G_S) \le \sqrt{n\left(2m + \sigma - \frac{\sigma^2}{n}\right)}.$$
 (6)

Proof. The expression

$$\sum_{i=1}^{n} \sum_{j=1}^{n} \left(\left| \lambda_i - \frac{\sigma}{n} \right| - \left| \lambda_j - \frac{\sigma}{n} \right| \right)^2$$

is evidently non-negative. Expanding it we get

$$n\sum_{i=1}^{n}\left|\lambda_{i}-\frac{\sigma}{n}\right|^{2}+n\sum_{j=1}^{n}\left|\lambda_{j}-\frac{\sigma}{n}\right|^{2}-2\sum_{i=1}^{n}\left|\lambda_{i}-\frac{\sigma}{n}\right|\sum_{j=1}^{n}\left|\lambda_{j}-\frac{\sigma}{n}\right|$$

which by Lemma 5 and Eq. (3) yields

$$2n\left(2m+\sigma-\frac{\sigma^2}{n}\right) - 2E(G_S)^2 \ge 0$$

from which Theorem 6 directly follows.

As expected, formula (6) reduces to the original McClelland bound [17] for $\sigma = 0$, but also for $\sigma = n$, in harmony with Lemma 1(b).

Acknowledgement. Izudin Redžepović and Boris Furtula were supported by the Serbian Ministry of Education, Science and Technological Development (Grant No. 451-03-9/2021-14/200122).

References

- J. Aihara, General rules for constructing Hückel molecular orbital characteristic polynomials, J. Am. Chem. Soc. 98 (1976) 6840–6844.
- [2] V. Consonni, R. Todeschini, New spectral index for molecule description, MATCH Commun. Math. Comput. Chem. 60 (2008) 3–14.
- [3] M. Cavers, S. Fallat, S. Kirkland, On the normalized Laplacian energy and general Randić index R₋₁ of graphs, *Lin. Algebra Appl.* **433** (2010) 172–190.
- [4] A. Graovac, I. Gutman, N. Trinajstić, T. Živković, Graph theory and molecular orbitals. Application of Sachs theorem, *Theor. Chim. Acta* 26 (1972) 67–78.
- [5] I. Gutman, Topological studies on heteroconjugated molecules. Alternant systems with one heteroatom, *Theor. Chim. Acta* 50 (1979) 287–297.
- [6] I. Gutman, Topological studies on heteroconjugated molecules. VI. Alternant systems with two heteroatoms, Z. Naturforsch. 45a (1990) 1085–1089.
- [7] I. Gutman, Rectifying a misbelief: Frank Harary's role in the discovery of the coefficient-theorem in chemical graph theory, J. Math. Chem. 16 (1994) 73–78.
- [8] I. Gutman, B. Furtula, The total π-electron energy saga, Croat. Chem. Acta 90 (2017) 359–368.
- [9] I. Gutman, B. Furtula, Energies of Graphs Survey, Census, Bibliography, Center Sci. Res., Kragujevac, 2019.
- [10] I. Gutman, H. Ramane, Research on graph energies in 2019, MATCH Commun. Math. Comput. Chem. 84 (2020) 277–292.
- [11] I. Gutman, B. Zhou, Laplacian energy of a graph, Lin. Algebra Appl. 414 (2006) 29–37.
- [12] F. Harary, *Graph Theory*, Addison–Wesley, Reading, 1969.
- [13] D. König, Theorie der endlichen und enendlichen Graphen, Akademische Verlagsgessellschaft, Leipzig, 1936.
- [14] X. Li, Y. Shi. I. Gutman, Graph Energy, Springer, New York, 2012.
- [15] R. B. Mallion, A. J. Schwenk, N. Trinajstić, A graphical study of heteroconjugated molecules, *Croat. Chem. Acta* 46 (1974) 171–182.
- [16] R. B. Mallion, N. Trinajstić, A. J. Schwenk, Graph theory in chemistry Generalization of Sachs' formula, Z. Naturforsch. 29a (1974) 1481–1484.
- [17] B. J. McClelland, Properties of the latent roots of a matrix: The estimation of π -electron energies, J. Chem. Phys. 54 (1971) 640–643.
- [18] H. S. Ramane, Energy of graphs, in: M. Pal, S. Samanta, A. Pal (Eds.), Handbook of Research on Advanced Applications of Graph Theory in Modern Society, IGI Global, Hershey, 2020, pp. 267–296.

- [19] M. J. Rigby, R. B. Mallion, A. C. Day, Comment on a graph-theoretical description of heteroconjugated molecules, *Chem. Phys. Lett.* **51** (1977) 178–182.
- [20] H. Sachs, Beziehungen zwischen den in einem Graphen enthaltenen Kreisen und seinem charakteristischen Polynom, Publ. Math. (Debrecen) 11 (1964) 119–134.
- [21] H. Sachs, Einführung in die Theorie der endlichen Graphen, Hauser, München, 1971.