

THE EFFECT OF INTERACTION OF SYSTEMATIC FACTORS ON MILK PERFORMANCE OF SIMMENTAL COWS IN STANDARD LACTATION

M.D. Petrović^{1*}, M.M. Petrović², Z. Skalicki³, V. Bogdanović³, R. Đoković¹, S. Rakonjac¹

¹Faculty of Agronomy, Cacak

²Institute for Animal Husbandry, Belgrade-Zemun

³Faculty of Agriculture, Belgrade-Zemun

*Corresponding author: Milun D. Petrovic, e-mail: milunp@tfc.kg.ac.rs

Original scientific paper

Abstract: The effect of interaction between the farm and calving season, as well as the farm and group of lactations on milk performance traits (milk yield, yield of milk fat, 4% fat corrected milk and milk fat content) was analyzed on sample of 2805 Simmental cows with total 9718 standard lactations, housed on dairy farm “Zlatiborski suvati” on Zlatibor (578 cows and 1968 lactations), dairy farm of the Agricultural holding Dobričevo in Čuprija (964 cows and 3237 lactations) and dairy animals housed on family farms/households in the region of Kotraž (1263 cows and 4513 lactations). The significance of the effect of interaction between the farm and calving season, as well as interaction between the farm and group of lactations on all milk performance traits was very high ($P < 0.001$), which justified their inclusion into models for assessment of the breeding value of dairy cows. The share of variance of interactions in total variance of milk performance traits in standard lactations was low (below and around 4%), which clearly indicated even more systematic factors and their interactions which influence the total variability of observed traits of milk performance of cows in standard lactations.

Key words: systematic factors, interaction, lactation, Simmental breed.

Introduction

The cattle breeding procedures used today most commonly employ linear methods and models that combine fixed parameters (year, farm, season, lactation) and random variables (age at first conception or calving, genetic effect of sire, genetic effect of an individual animal, etc.) which can be mutually dependent (related) or independent, with or without interactions, depending on the trait analysed. The selected model results essentially in the breeding value of an individual animal (Bogdanović *et al.*, 2003).

Depending on the level of production, sample size and mathematical statistical model, non-genetic discontinuous factors (year, farm, season, lactation) can account for as much as above 50% of the total variations in milk production (*Stojić et al., 1996*). The same authors, *Stojić et al., 1995*, reported that farm, year and season of calving induced 35.7% of total variations in milk production. A somewhat lower percentage of non-genetic factors (farm, year, calving season and lactation number) of 22.7% in the total variability of performance traits was found by *Jovanovac (1987)*, whereas *Hansen et al., 1983* reported a considerably higher percentage of about 45% (in terms of the farm-year-season effect).

The effect of season of birth of cows and their calving, i.e. beginning of lactation as systematic factor, on traits of milk performance is reflected through various climatic circumstances and nutrition throughout the year, so it is included into models for evaluation of breeding value of dairy animals.

Moat of domestic and foreign authors indicate significant and highly significant effect of order of lactation, i.e. groups of lactations on expression of production traits. Namely, maximum production of milk is achieved depending on the intensity of breeding in the period from the third until the fifth lactation. The lowest production of milk is in the first lactation, due to insufficient body development of animals, whereas drop in production occurring after the third i.e. fifth lactation often occurs due to health disorders.

Breeding region or farm usually have significant impact on milk performance traits due to different rearing methods, nutrition, housing, care, climatic conditions, age structure and herd size, as well as series of other influences related to work and farm management.

Material and methods

The effect of systematic, non-genetic factors and their interactions on traits of milk performance in standard lactations was analyzed on sample of 2805 Simmental cows with total 9718 standard lactations, housed on dairy farm "Zlatiborski suvati" on Zlatibor, dairy farm of the Agricultural holding Dobričevo in Čuprija and dairy animals housed on family farms/households in the region of Kotraž.

Distribution of standard lactations according to classes of systematic influences is presented in Table 1.

Table 1. Presentation of data across different classes of major systematic effects

Farm	Lactat. No.	Cow no.	Lactat. group	Lactat. No.	Year of birth	Lactat. No.	Season of birth	Lactat. No.	S. of calving	Lactat. No.
I (Zlatibor)	1968	578	I (1)	2800	1982.	847	I	2298	I	2281
			II (2)	2297	1983.	565	II	2837	II	2642
II (Dobričevo)	3237	964	III (3)	1715	1984.	624	III	2035	III	2269
			IV(4)	1213	1985.	541	IV	2552	IV	2526
III (Kotražā)	4513	1263	V (5)	799	1986.	855				
			VI(6+7+8+9+10+11+12)	894	1987.	822				
					1988.	791				
					1989.	839				
					1990.	786				
					1991.	627				
					1992.	612				
					1993.	306				
					1994.	467				
					1995.	516				
					1996.	342				
		1997.	76							
		1998.	102							

The impact of following systematic factors on expression of production traits in standard lactations, such as milk yield in standard lactation (MYSL, kg), milk fat content in standard lactations (MFCSL, %), milk fat yield in standard lactations (MFYSL, kg) and yield of 4 % fat corrected milk in standard lactations (4%FCMSL, kg), was observed:

- *Breeding area.* The effect of three farm locations was studied (the farm on Mt. Zlatibor, Dobričevo farm and farms in the Kotražā region).
- *Lactation groups.* Lactation groups were established in order to equalise the number of animals within different lactations as much as possible and reduce variability:
 - Group I (first lactation),
 - Group II (second lactation),
 - Group III (third lactation),
 - Group IV (fourth lactation),
 - Group V (fifth lactation),
 - Group VI (sixth and other lactations),
- *Calving season,* i.e. the onset of lactation (I-spring season (March through May), II-summer season (June through August), III-autumn season (September through November), IV-winter season (December through February)).
- *Year of birth x season of birth interaction* (cows that calved from 1982 to 1998 x 4 seasons (I-spring, II-summer, III-autumn and IV-winter seasons)).

- *Breeding area x calving season interaction* (3 breeding areas x 4 calving seasons).
- *Breeding area x lactation group interaction* (3 breeding areas x 6 lactation groups).
- *Age at first conception*.

For analysis of the mentioned non-genetic factors and their interactions on production traits in standard lactations, a general linear model was used based on application of the method of least squares of the statistical software Statistica for Windows Release 6.0, which in this case is:

$$y_{ijkl} = \mu + B_i + L_j + G_k + Cs_l + BCs_{il} + BL_{ij} + b_1(x_{i1} - \bar{x}_1) + e_{ijkl}, \text{ where:}$$

y_{ijkl} – an individual animal of the i -th breeding area, j -th lactation group, k -th group, l -th calving season,

μ – population mean with equal participation of all classes of effects (B, L, G, Cs, BCs, BL),

B_i – fixed effect of the i -th breeding area (1-3),

L_j – fixed effect of the j -th lactation group (1-6),

G_k – fixed effect of the k -th group (year of birth x season of birth) (1-68),

Cs_l – fixed effect of the l -th calving season (1-4),

BCs_{il} – fixed effect of the i -th breeding area x l -th calving season interaction (1-12),

BL_{ij} – fixed effect of the i -th breeding area x j -th lactation group interaction (1-18),

b_1 – linear regression coefficient of the effect of age at first conception, and

e_{ijkl} – other undetermined effects.

Results and discussion

In research results only the results of the effect of interaction between systematic factors on expression of milk performance traits in standard lactations according to applied model are presented.

Mean values of least squares, standard errors and significance of the influence of interaction farm and calving season on production traits of standard lactations according to applied model are presented in Table 2.

Table 2. Mean values of least squares, standard errors and significance of the influence of interaction farm and calving season on production traits of standard lactations

Systematic factors			MYSL (kg)		MFCSL (%)		MFYSL (kg)		4%FCMSL (kg)	
Farm	Calving season	N	LSM	SE _{LSM}	LSM	SE _{LSM}	LSM	SE _{LSM}	LSM	SE _{LSM}
I	I	374	4460,4	41,734	3,66	0,008	162,79	1,582	4226,0	40,220
I	II	425	4245,2	39,206	3,67	0,007	155,21	1,486	4026,2	37,783
I	III	593	4320,1	33,228	3,68	0,006	158,64	1,260	4107,6	32,022
I	IV	576	4394,5	33,430	3,67	0,006	160,76	1,267	4169,2	32,217
II	I	705	4007,8	31,350	3,79	0,006	151,64	1,188	3877,7	30,212
II	II	771	3860,7	30,003	3,77	0,005	145,50	1,137	3726,8	28,914
II	III	908	4168,7	28,327	3,78	0,005	157,60	1,074	4031,5	27,299
II	IV	853	4182,3	29,103	3,80	0,005	158,85	1,103	4055,7	28,047
III	I	1202	3962,4	24,988	3,91	0,005	154,89	0,947	3908,4	24,081
III	II	1446	3925,9	22,548	3,91	0,004	153,60	0,855	3874,4	21,729
III	III	768	3962,3	29,093	3,91	0,005	155,02	1,103	3910,2	28,037
III	IV	1097	4005,9	25,476	3,91	0,005	156,61	0,966	3951,5	24,551
F _{exp}			9.39 ^{***}		4.60 ^{***}		9.82 ^{***}		9.71 ^{***}	

N.S. - $P > 0.05$; * - $P < 0.05$; ** - $P < 0.01$; *** - $P < 0.001$;

The effect of interaction between the farm and calving season on all milk traits of standard lactations was highly significant ($P < 0.001$), therefore, inclusion not only of these individual systematic factors but also their interactions into models for evaluation of breeding value of cows is justified. The effect farms-year-season, according to research by *Stojić (1996)*, was the most dominant effect for all studied milk performance indicators in standard lactation ($P < 0.01$). *Petrović M.M. et al., 1997* and *Petrović D.M. et al. 2009*, in their study of the active population of Simmental cattle in Serbia, indicated that interaction between breeding region and calving season contributed to significant variation in the milk yield and yield of milk fat ($P < 0.01$). Very significant ($P < 0.01$) and significant ($P < 0.05$) effect of breeding region, as individual systematic factor, on production traits of Simmental cows is indicated by *Petrović et al., 1997 and 2006*, *Perišić, 1998*, *Rychen, 1999*, *Chladek and Kucera, 2000*, *Đurđević, 2001*, *Đurđević et al., 2002*, *Panić, 2005* and *Petrović D.M. et al., 2009*.

Share of interaction between farm and calving season in total variance of milk traits in standard lactations was very low and ranged from only 0.18% in case of milk fat yield to 0.49% in case of production of 4% fat corrected milk, which clearly indicated the high contribution of other systematic factors in variability of these traits which are included into presented model and were not considered in this study. Low level of share of variance of certain systematic factors in total variability of production of 4% fat corrected milk is stated in the research of *Petrović D.M. et al., 2010*.

Results of the analysis of the effect of interaction between breeding region and groups of lactation on production traits of standard lactations, i.e. means of least squares (LSM), standard errors of means (SE_{LSM}) and significance of observed effects are presented in Table 3.

Table 3. Means of least squares, standard errors of means and significance of the effect of interaction between farm and groups of lactation on production traits of standard lactations

Systematic factors			MYSL (kg)		MFCSL (%)		MFYSL (kg)		4%FCMSL (kg)	
Farm	Gr.lac.	N	LSM	SE_{LSM}	LSM	SE_{LSM}	LSM	SE_{LSM}	LSM	SE_{LSM}
I	I	574	3941,7	31,915	3,70	0,006	144,82	1,210	3749,0	30,757
I	II	472	4384,6	35,782	3,65	0,007	159,56	1,357	4147,2	34,484
I	III	336	4563,3	42,249	3,65	0,008	166,13	1,602	4317,3	40,716
I	IV	233	4548,4	50,146	3,66	0,009	166,17	1,901	4311,9	48,326
I	V	172	4489,0	58,154	3,66	0,011	163,91	2,205	4254,2	56,044
I	VI	181	4203,2	56,921	3,71	0,010	155,50	2,158	4013,7	54,856
II	I	964	3916,1	26,285	3,81	0,005	149,20	0,996	3804,5	25,331
II	II	793	4167,7	28,559	3,80	0,005	158,18	1,083	4039,7	27,522
II	III	593	4179,5	32,534	3,79	0,006	158,03	1,233	4042,2	31,353
II	IV	402	4154,0	38,776	3,77	0,007	156,40	1,470	4007,6	37,369
II	V	240	4044,7	49,276	3,77	0,009	152,48	1,868	3905,2	47,488
II	VI	245	3867,5	49,082	3,78	0,009	146,09	1,861	3738,4	47,300
III	I	1262	3313,3	23,349	3,94	0,004	130,80	0,885	3287,3	22,501
III	II	1032	3633,3	25,211	3,93	0,005	142,69	0,956	3593,6	24,296
III	III	786	3970,1	28,574	3,91	0,005	155,18	1,083	3915,7	27,537
III	IV	578	4208,3	32,819	3,90	0,006	163,96	1,244	4142,8	31,628
III	V	387	4338,5	39,479	3,90	0,007	169,04	1,497	4271,0	38,046
III	VI	468	4321,1	36,825	3,90	0,007	168,52	1,396	4256,2	35,488
F_{exp}			49.41***		4.10***		49.87***		50.14***	

N.S. - $P > 0.05$; * - $P < 0.05$; ** - $P < 0.01$; *** - $P < 0.001$;

Significance of the effect of interaction between breeding region and group of lactations on all milk traits of standard lactations was highly significant ($P < 0.001$), which justified its inclusion into evaluation of breeding value of dairy animals, as confirmed by research of *M.M. et al., (1997)* and *Petrović D.M. et al. (2009)*.

Share of variance of interaction between breeding region and group of lactations in total variance of production traits of whole lactations was slightly higher and ranged from 0.25% in milk fat content to 4.12% in milk fat yield in standard lactations. However, the share of variance in total variability

unambiguously indicates even more systematic factors and their interactions which influence total variability of observed milk traits of standard lactations, as confirmed by *Petrović M.M. et al., (1997)* and *Petrović D.M. et al. (2009 and 2010)*.

Conclusion

Based on presented results the following can be concluded:

- The significance of the effect of interaction between farm and calving season, as well as interaction between farm and group of lactations on all milk traits of standard lactations was very high ($P < 0.001$), which justifies their inclusion into models for evaluation of breeding value of dairy cows.
- The share of variance of interactions in total variance of milk traits of standard lactations was low, which clearly indicated presence of more systematic factors and their interactions which have influence on total variability of observed milk traits of standard lactations.

Acknowledgment

This work was financed by the Ministry of Education and Science, Republic of Serbia, projects TR 31086 and TR 31053.

Efekat interakcije sistematskih faktora na mlečnost krava simentalske rase u standardnim laktacijama

M.D. Petrović, M.M. Petrović, Z. Skalicki, V. Bogdanović, R. Đoković, S. Rakonjac

Rezime

Uticaj interakcije farme i sezone telenja kao i farme i grupe laktacija na osobine mlečnosti (proizvodnja mleka, mlečne masti, 4% mast-korigovanog mleka i sadržaj mlečne masti) analizirano je kod 2805 krava simentalske rase sa ukupno 9718 standardnih laktacija koje su bile smeštene na farmi muznih krava "Zlatiborski suvati" na Zlatiboru (578 krava i 1968 laktacija), farmi muznih krava na Poljoprivrednom dobru Dobričevo u Čupriji (964 krava i 3237 laktacija) i području Kotraže na individualnim gazdinstvima (1263 krava i 4513 laktacija).

Značajnost uticaja interakcija farme i sezone telenja, kao i interakcije farme i grupe laktacija na sve osobine mlečnosti standardnih laktacija bila je vrlo

visoka ($P < 0.001$), što opravdava njihovo uključivanje u modele za ocenu priplodne vrednosti muznih krava.

Udeo varijanse interakcija u ukupnoj varijansi osobina mlečnosti standardnih laktacija bio je nizak (ispod i oko 4%), što jasno ukazuje na još veliki broj sistematskih faktora i njihovih interakcija koji utiču na ukupnu varijabilnost posmatranih osobina mlečnosti krava u standardnim laktacijama.

References

- BOGDANOVIĆ V., ĐEDOVIĆ RADICA (2003): Testiranje i metode procene priplodne vrednosti simentalčkih bikova. Savremeni trendovi u mlekarnstvu, Zbornik radova, str. 46-50. Zlatibor.
- CHLADEK G., KUCERA J. (2000): An analysis of some factors affecting the milk production of cows sired by Montbeliard sires in the Czech Republic. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*. 2000, 48: 5, 21-26; 10 ref.
- ĐURĐEVIĆ R. (2001): Genetička analiza mlečnosti i reprodukcijskih svojstava krava simentalčke rase. Doktorska disertacija. Poljoprivredni fakultet Novi Sad, 2001.
- ĐURĐEVIĆ R., VIDOVIĆ V., ANTOV G., LATINOVIĆ D. (2002): Genetička varijabilnost perzistencije laktacije krava simentalčke rase. *Biotechnology in Animal Husbandry* 18(5-6), p.9-15.
- HANSEN L., FREEMAN A.E., BERGER P.J. (1983): Variances, Repeatabilities and Age Adjustments of Yield and Fertility in Dairy Cattle. *Journal of Dairy Science*, 66, 281-292.
- PANIĆ JOVANKA (2005): Kvantitativno – genetička analiza svojstava mlečnosti krava simentalčke rase. Poljoprivredni fakultet, Univerzitet u Novom Sadu (Magistarska teza).
- PERIŠIĆ P. (1998): Reproktivne i proizvodne osobine različitih genotipova krava simentalčke rase. Magistarska teza, Poljoprivredni fakultet, Beograd.
- PETROVIĆ D.M., BOGDANOVIĆ V., PETROVIĆ M.M., SNEŽANA BOGOSAVLJEVIĆ-BOŠKOVIĆ (2010): Uticaj paragenetskih faktora na proizvodnju 4% mast-korigovanog mleka u celim i standardnim laktacijama. XV savetovanje o biotehnologiji. Zbornik radova Vol.15(17), str. 585-590, 2010. Agronomski fakultet, Čačak.
- PETROVIĆ D.M., SKALICKI Z., PETROVIĆ M.M., BOGDANOVIĆ V. (2009): The Effect of Systematic Factors on Milk Yield in Simmental Cows Over Complete Lactations. *Biotechnology in Animal Husbandry* 25(1-2), p 61-71, Belgrade.

- Petrović M.M., Lazarević R., Lazarević L.J., Aleksić S., Mišćević B., Perković S. (1997): Proizvodni efekti selekcije aktivne populacije simentalских goveda u Srbiji. *Bootehnologija u stočarstvu*, god. 13, Br. 3-4, str. 57-64, Beograd.
- PETROVIĆ M.M., SRETENOVIĆ LJILJANA, PANTELIĆ V., ALEKSIĆ S., MIŠĆEVIĆ B., BOGDANOVIĆ V., OSTOJIĆ DUŠICA, PETROVIĆ D.M. (2006): Results of the Application of the Technology of Genetic Improvement of Simmental Cattle Population in Serbia. *Biotechnology in Animal Husbandry* 22(1-2), p 1-8, Belgrade.
- RYCHEN M. (1999): A 6279 kg milk yield – where are the limits? *Schweizer-Fleckvieh*. 1999, No. 7, 26-39.
- SONJA JOVANOVAČ (1987): Utjecaj sistematskih faktora okoline na mliječnost krava Holstein-Friesian pasmine. *Znanost i praksa u poljoprivrednoj i prehrambenoj tehnologiji* 17(3-4), 303-314.
- STATSOFT INC. STATISTICA FOR WINDOWS, Version 6.0, Computer program manual. Tulsa, : StatSoft Inc., 1995.
- STOJIĆ P., KATIĆ M., LAZAREVIĆ L.J., LATINOVIĆ D., TRIFUNOVIĆ G., RADMILA BESKOROVAJNI, BRKIĆ N. (1995): Ponovljivost dnevnih prinosa mleka tokom laktacije crno belih krava. Prvi Simpozijum za oplemenjivanje organizama sa međunarodnim učešćem. Vrnjačka Banja.
- STOJIĆ P. (1996): Faktori korekcije osobina mlečnosti i njihov doprinos oceni priplodne vrednosti bikova i krava. Poljoprivredni fakultet, Univerzitet u Beogradu, Doktorska disertacija.