



University of Belgrade
Faculty of Mechanical Engineering

Numerical Methods for Large Scale Problems

*Dedicated to Professor Lothar Reichel
(Kent State University, Ohio, USA)
on the occasion of his 70th Anniversary*

A B S T R A C T B O O K



Belgrade, June 6–10, 2022

Scientific committee:

- Alessandro Buccini (University of Cagliari, Italy)
- Stefano Cipolla (University of Padua, Italy)
- Maria Carmela De Bonis (University of Basilicata, Italy)
- Marco Donatelli (University of Insubria, Como, Italy)
- Christos Koukouvinos (National Technical University of Athens, Greece)
- Gradimir Milovanović (Serbian Academy of Sciences and Arts, Serbia)
- Donatella Occorsio (University of Basilicata, Italy)
- Mirjeta Pasha (Arizona State University, USA)
- Michela Redivo-Zaglia (University of Padova, Italy)
- Lothar Reichel (Kent State University, OHIO, USA)
- Giuseppe Rodriguez (University of Cagliari, Italy)
- Hassane Sadok (Université du Littoral Côte d'Opale, France)
- Miodrag Spalević (University of Belgrade, Serbia)
- Marija Stanić (University of Kragujevac, Serbia)
- Raf Vandebril (KU Leuven, Belgium)

Organizing committee:

- Alessandro Buccini (University of Cagliari, Italy)
- Dušan Djukić (University of Belgrade, Serbia)
- Marco Donatelli (University of Insubria, Como, Italy)
- Laura Dykes (University School, U.S.)
- Luisa Fermo (University of Cagliari, Italy)
- Davorika Jandrlić (University of Belgrade, Serbia)
- Rada Mutavdžić Djukić (University of Belgrade, Serbia)
- Mirjeta Pasha (Arizona State University, U.S.)
- Aleksandar Pejčev (University of Belgrade, Serbia)
- Giuseppe Rodriguez (University of Cagliari, Italy), chair
- Miodrag Spalević (University of Belgrade, Serbia), chair
- Jelena Tomanović (University of Belgrade, Serbia)

Anti-Gaussian quadrature rules for the optimal set of quadrature rules in Borges' sense

Nevena Z. Petrović¹, Marija P. Stanić¹, and Tatjana V. Tomović Mladenović¹

Department of Mathematics and Informatics, Faculty of Science, University of Kragujevac
Radoja Domanovića 12, 34000 Kragujevac, Serbia
`nevena.z.petrovic,marija.stanic,tatjana.tomovic@pmf.kg.ac.rs`

Abstract

Anti-Gaussian quadrature rules, introduced by Laurie in [1], have the property that their error is equal in magnitude but of the opposite sign to the corresponding Gaussian quadrature rules. Guided by that idea, we define and analyse anti-Gaussian quadrature rules for the optimal set of quadrature rules in Borges' sense (see [2]), with respect to the set of r different weight functions. Also, we introduce the set of averaged quadrature rules and give some numerical examples.

Keywords: Anti-Gaussian quadratures, Optimal set of quadrature rules in Borges' sense, Weight function

References

1. D.P. LAURIE, *Anti-Gaussian quadrature formulas*, Math. Comp. 65(214), (1996) 739-747.
2. C.F. BORGES, *On a class of Gauss-like quadrature rules*, Numer. Math. 67, (1994) 271-288.