



Relationships between Some Distance–Based Topological Indices

Hongbo Hua^a, Ivan Gutman^b, Hongzhuan Wang^a, Kinkar Ch. Das^c

^aFaculty of Mathematics and Physics, Huaiyin Institute of Technology,
Huai'an, Jiangsu 223003, PR China

^bFaculty of Science, University of Kragujevac,
34000 Kragujevac, Serbia

^cDepartment of Mathematics, Sungkyunkwan University,
Suwon 440-746, Republic of Korea

Abstract. The Harary index (HI), the average distance (AD), the Wiener polarity index (WPI) and the connective eccentricity index (CEI) are distance–based graph invariants, some of which found applications in chemistry. We investigate the relationship between HI , AD , and CEI , and between WPI , AD , and CEI . First, we prove that $HI > AD$ for any connected graph and that $HI > CEI$ for trees, with only three exceptions. We compare WPI with CEI for trees, and give a classification of trees for which $CEI \geq WPI$ or $CEI < WPI$. Furthermore, we prove that for trees, $WPI > AD$, with only three exceptions.

1. Introduction

Throughout this paper we consider only simple connected graphs. For a graph $G = (V, E)$ with vertex set $V = V(G)$ and edge set $E = E(G)$, the *degree* of a vertex v , denoted by $d_G(v)$, is the number of edges incident with v . Denote by $d_G(u, v)$ the distance between vertices u and v in G . The *eccentricity* of a vertex v in a graph G is defined to be $\varepsilon_G(v) = \max\{d_G(u, v) | u \in V(G)\}$. The *diameter* of a connected graph G is equal to $\max\{\varepsilon_G(v) | v \in V(G)\}$, whereas its *radius* is equal to $\min\{\varepsilon_G(v) | v \in V(G)\}$.

A connected graph is said to be a *tree* if it contains no cycles. Let P_n , S_n , C_n , and K_n be the path, star, cycle, and complete graph of order n , respectively. For other notation and terminology not defined here, the readers are referred to [3].

One of the oldest and best studied distance–based graph invariants is the Wiener index, defined as [26]

$$W(G) = \sum_{\{u,v\} \subseteq V(G)} d_G(u, v).$$

2010 *Mathematics Subject Classification.* 05C35; 05C12

Keywords. distance (in graph); Harary index; average distance; Wiener polarity index; connective eccentricity index

Received: 08 February 2018; Accepted: 15 October 2018

Communicated by Francesco Belardo

Hua was supported by National Natural Science Foundation of China under Grant No. 11571135 and Qing Lan Project of Jiangsu Province, P.R. China. Wang was supported by National Natural Science Foundation of China under Grant No. 11571135. Das was supported by the Sungkyun research fund, Sungkyunkwan University, and National Research Foundation of the Korean government with grant No. 2017R1D1A1B03028642.

Email addresses: hongbo_hua@163.com (Hongbo Hua), gutman@kg.ac.rs (Ivan Gutman), wanghz412@163.com (Hongzhuan Wang), kinkardas2003@gmail.com (Kinkar Ch. Das)

In some applications, it is more convenient to study the *average distance* (AD) of G ,

$$\overline{W}(G) = \frac{1}{\binom{n}{2}} \sum_{\{u,v\} \subseteq V(G)} d_G(u,v) = \frac{2}{n(n-1)} W(G).$$

Results on Wiener index can be found in the reviews [12, 16, 27]. For results on average distance see [4–6, 11] and the references cited therein.

Another distance-based graph invariant, put forward independently in [22] and [25], is the reciprocal-analogue of the Wiener index, named *Harary index* (HI), and defined as

$$H(G) = \sum_{\{u,v\} \subseteq V(G)} \frac{1}{d_G(u,v)}.$$

The *Wiener polarity index* (WPI), introduced also by Wiener in 1947 [26], is

$$W_p(G) = |\{(u,v) \mid d_G(u,v) = 3, u, v \in V(G)\}|.$$

It also found applications in chemistry [18, 24]. For recent mathematical results on WPI see [13–15, 21, 23, 30].

In 2000, the *connective eccentricity index* (CEI) of a connected graph G , denoted by $C^\xi(G)$, was introduced by Gupta et al. [17] as

$$C^\xi(G) = \sum_{u \in V(G)} \frac{d_G(u)}{\varepsilon_G(u)}.$$

For recent results on the CEI see [1, 28, 29] and the references cited therein.

Relationships between various graph invariants have received much attention over the past few decades, see e.g., [7–10, 19, 20].

In this paper, we investigate various relationships between the above listed distance-based graph invariants. We prove that the Harary index is greater than the average distance for any connected graph. Also, we prove that for trees, the Harary index is greater than the connective eccentricity index, with only three exceptions. Moreover, we compare the Wiener polarity index with the connective eccentricity index for trees, and give an explicit classification of all trees for which CEI is greater or smaller than WPI. We prove that for trees, the Wiener polarity index is greater than the average distance, with only three exceptions. Finally, we compare the Harary index with connective eccentricity index in terms of a radius-dependent condition.

2. Main Results

In this section, we investigate the relationship between the Harary index and average distance and connective eccentricity index, and the relationship between the Wiener polarity index and average distance and connective eccentricity index. We will proceed by dividing our discussions into four subsections.

2.1. Harary index and average distance

For a connected graph G , the *remoteness* of G is defined as $\rho = \rho(G) = \max_{v \in V(G)} \frac{1}{n-1} D_G(v)$. We need a result on remoteness due to Aouchiche and Hansen, which reads as follows:

Lemma 2.1 ([2]). *Let G be a connected graph of order n with remoteness ρ . Then $\rho \leq n/2$ with equality if and only if $G \cong P_n$.*

Next, we will show that the Harary index is greater than the average distance for any connected graph. First, we prove a somewhat stronger result:

Theorem 2.2. Let G be a connected graph with average distance $\overline{W}(G)$ and average degree $\overline{d}(G)$. Then

$$H(G) > \overline{d}(G) \cdot \overline{W}(G).$$

Proof. Suppose that the order and size of G are n and m , respectively. Then $\overline{d}(G) = \frac{2m}{n}$. By Lemma 2.1,

$$\overline{d}(G) \cdot \overline{W}(G) = \frac{2m}{n} \cdot \frac{\sum_{v \in V(G)} D_G(v)}{n(n-1)} = \frac{2m}{n^2} \cdot \sum_{v \in V(G)} \frac{D_G(v)}{n-1} \leq \frac{2m}{n^2} \cdot n\rho \leq \frac{2m}{n^2} \cdot n \cdot \frac{n}{2} = m.$$

Obviously,

$$H(G) = \sum_{\{u,v\} \subseteq V(G)} \frac{1}{d_G(u,v)} \geq m$$

with equality if and only if $G \cong K_n$.

Therefore, $H(G) \geq \overline{d}(G) \cdot \overline{W}(G)$. It is not difficult to see that the equality in the above inequality cannot be attained. Thus, $H(G) > \overline{d}(G) \cdot \overline{W}(G)$. \square

Since $\overline{d}(G) \geq 1$ for any connected graph G , we have:

Corollary 2.3. Let G be a connected graph with average distance $\overline{W}(G)$ and Harary index $H(G)$. Then

$$H(G) > \overline{W}(G).$$

2.2. Harary index and connective eccentricity index

In order to find the relationship between the Harary index and the connective eccentricity index, we first consider the following three special graphs.

For the complete graph K_n , $C^\xi(K_n) = n(n-1) > \frac{n(n-1)}{2} = H(K_n)$ for $n \geq 2$.

For $a \geq 1, b \geq 1$, let S_{a+1} and S_{b+1} be stars on $a+1$ and $b+1$ vertices, respectively. Then the double star $S_{a,b}$ is the tree obtained by connecting an edge between two centers of S_{a+1} and S_{b+1} .

For the double star $S_{a,b}$ ($a+b = n-2$), $C^\xi(S_{a,b}) = \frac{5n-4}{6}$, $H(S_{a,b}) = \frac{ab}{3} + \frac{3(a+b)}{2} + 1 \geq \frac{n-3}{3} + \frac{3(n-2)}{2} + 1 = \frac{11n-18}{6}$. Therefore, $H(S_{a,b}) > C^\xi(S_{a,b})$ for $n \geq 4$.

For the complete bipartite graph $K_{\frac{n}{2}, \frac{n}{2}}$, where $n \geq 4$ and n is an even integer, $C^\xi(K_{\frac{n}{2}, \frac{n}{2}}) = \frac{n^2}{4}$, $H(K_{\frac{n}{2}, \frac{n}{2}}) = \frac{3n^2}{8} - \frac{n}{4}$. Therefore, $C^\xi(K_{\frac{n}{2}, \frac{n}{2}}) < H(K_{\frac{n}{2}, \frac{n}{2}})$ for $n \geq 4$.

From the above examples, one concludes that in the general case, HI and CEI are incomparable. Bearing this in mind, we shall restrict our considerations to trees.

Theorem 2.4. Let T be a tree of order n . If $T \in \{P_2, P_3\}$, then $H(T) < C^\xi(T)$. Otherwise,

$$H(T) \geq C^\xi(T)$$

with equality if and only if $T \cong S_4$.

Proof. For $T \in \{P_2, P_3\}$, it can be easily checked that $H(T) < C^\xi(T)$. Assume therefore that $T \notin \{P_2, P_3\}$. Then, $n \geq 4$.

Let $A = \{v | d_T(v) = n-1\}$. Since T is a tree, we have $|A| \leq 1$.

If $|A| = 0$, then $\varepsilon_T(v) \geq 2$ for each vertex v in T , and thus, $\frac{d_T(v)}{\varepsilon_T(v)} \leq \frac{d_T(v)}{2}$. For any vertex v in T , write $\widehat{D}_T(v) = \sum_{u \in V(G) \setminus \{v\}} \frac{1}{d_T(u,v)}$. Note that if $\varepsilon_T(v) \geq 2$, then $\widehat{D}_T(v) > d_T(v)$. Therefore,

$$H(T) = \frac{1}{2} \sum_{v \in V(T)} \widehat{D}_T(v) > \frac{1}{2} \sum_{v \in V(T)} d_T(v) \geq \sum_{v \in V(T)} \frac{d_T(v)}{\varepsilon_T(v)} = C^\xi(T).$$

Thus, in this case, $H(T) > C^\xi(T)$.

Now, let $|A| = 1$. Then $T \cong S_n$, and thus, $H(T) = H(S_n) = (n-1) + \binom{n-1}{2} \cdot \frac{1}{2} = \frac{(n-1)(n+2)}{4}$, $C^\xi(T) = \frac{n-1}{2} + (n-1) = \frac{3(n-1)}{2}$. Note that $n \geq 4$. Thus, $H(T) \geq C^\xi(T)$, with equality only if $n = 4$, that is, if $T \cong S_4$. Conversely, if $T \cong S_4$, then $H(T) = C^\xi(T)$.

This completes the proof. \square

Next, we compare the Harary and connective eccentricity indices for connected graphs under given restricted condition.

Theorem 2.5. Let G be a connected graph of order $n \geq 4$ with m edges and $p \geq 0$ vertices of degree $n - 1$. If $m \geq \frac{(n-1)(n-2p)}{2}$ and $p < \frac{n}{2}$, then $H(G) \leq C^\xi(G)$.

Proof. Let Δ be the maximum degree of graph G . First, we claim that $\Delta = n - 1$.

If $\Delta \leq n - 2$, then $p = 0$. By our assumption that $m \geq \frac{(n-1)(n-2p)}{2} = \frac{n(n-1)}{2}$, we must have $G \cong K_n$, a contradiction.

Thus, we may assume that $\Delta = n - 1$. Then $d \leq 2$. If $d = 1$, then $G \cong K_n$ and hence $H(G) = \frac{n(n-1)}{2} < n(n-1) = C^\xi(G)$. Suppose therefore that $d = 2$. Since $m \geq \frac{(n-1)(n-2p)}{2}$, we have

$$H(G) = m + \frac{1}{2} \left[\frac{n(n-1)}{2} - m \right] \leq p(n-1) + \frac{2m - p(n-1)}{2} = C^\xi(G).$$

\square

2.3. Wiener polarity index and connective eccentricity index

In order to find a relationship between the Wiener polarity index and the connective eccentricity index, we first consider the following two special graphs.

For the complete graph K_n , $C^\xi(K_n) = n(n-1) > 0 = W_p(K_n)$ for $n \geq 2$.

For the path P_n , $C^\xi(P_n) < n \cdot \frac{2}{2} = 4 < n - 3 = W_p(P_n)$ for $n \geq 8$.

These examples imply that in the general case, WPI and CEI are incomparable. In view of this, we restrict our considerations to trees.

We first introduce a special class of trees.

Suppose that $P_{d+1} = v_0v_1 \cdots v_{d-1}v_d$ is a path of length d . For $d \geq 3$, let $T_n(r, t)$ be a tree of order n with diameter d obtained from P_{d+1} by attaching r and t pendent vertices to v_1 and v_{d-1} , respectively. Here, $r \geq 0$, $t \geq 0$, and $r + t = n - d - 1$, See Fig. 1.




Fig. 1. The tree $T_n(r, t)$, where $r \geq 0$, $t \geq 0$, and $r + t = n - d - 1$.

We first state a result due to Deng et al.

Lemma 2.6 ([14]). Let T be a tree of order n and diameter $d \geq 3$. Then $W_p(T) \geq n - 3$ with equality if and only if $T \cong T_n(r, t)$ for $d > 4$, and $T \cong T_n(n - 4, 0)$ for $d = 3$.

Note that the tree $T_n(n - 4, 0)$ in Lemma 2.6 is isomorphic to the double star $S_{1, n-3}$.

In 2009, Du et al. gave the following remarkable formula for computing the Wiener polarity index of trees.

Lemma 2.7 ([15]). Let T be a tree. Then

$$W_p(T) = \sum_{uv \in V(T)} [d_T(u) - 1][d_T(v) - 1]. \tag{1}$$

Theorem 2.8. Let T be a tree of order n . If $T \in \{S_n, P_5, S_{2,2}, T_6(1, 0), T_7(2, 0)\}$ or $T \cong S_{1, n-3}$ for $4 \leq n \leq 14$, then $C^\xi(T) \geq W_p(T)$ with equality if and only if $T \cong T_7(2, 0)$ or $T = \text{cong}S_{1,11}$. Otherwise,

$$W_p(T) > C^\xi(T).$$

Proof. We have to separately consider the following three cases:

Case $d = 2$: Then $T \cong S_n$ and thus $C^\xi(T) = 3(n - 1)/2 > 0 = W_p(T)$.

Case $d = 3$: Then $T \cong S_{a,b}$ ($a + b + 2 = n$, $1 \leq a \leq b$) and thus

$$W_p(T) - C^\xi(T) = ab - \frac{5a + 5b + 6}{6} = \frac{5b(a - 1) + a(b - 5) - 6}{6}.$$

First we assume that $a = 1$. Then $T \cong S_{1, n-3}$ and then $W_p(T) - C^\xi(T) = \frac{n-14}{6}$. For $4 \leq n \leq 13$, $W_p(T) < C^\xi(T)$, for $n = 14$, $W_p(T) = C^\xi(T)$, whereas for $n \geq 15$, $W_p(T) > C^\xi(T)$. Assume next that $a \geq 2$. If $b = 2$, then $(a, b) = (2, 2)$ and therefore $W_p(T) < C^\xi(T)$. Otherwise, $b \geq 3$. Then $5b(a - 1) + a(b - 5) - 6 \geq 5b - 2a - 6 > 0$. Thus we have $W_p(T) > C^\xi(T)$.

Case $d \geq 4$: Assume first that $T \cong T_n(r, t)$. Then $W_p(T) \geq n - 3$, $C^\xi(G) \leq \frac{r+s+2}{4} + \frac{2n-6-r-s}{3} + \frac{2}{2}$, and

$$W_p(T) - C^\xi(T) \geq \frac{n}{3} - \frac{5}{2} + \frac{r+s}{12}.$$

If $n \geq 8$, then from the above it follows $W_p(T) > C^\xi(T)$. Otherwise, in this case $n = 5$ or 6 or 7 . Since $T \cong T_n(r, t)$, we have $T \cong P_5$ ($n = 5$) or $T \cong P_6$ ($n = 6$) or $T \cong T_6(1, 0)$ ($n = 6$) or $T \cong P_7$ ($n = 7$) or $T \cong T_7(1, 0)$ ($n = 7$) or $T \cong T_7(2, 0)$ ($n = 7$) or $T \cong T_7(1, 1)$ ($n = 7$).

For $T \cong P_5$, $C^\xi(T) = 17/6 > 2 = W_p(T)$. For $T \cong P_6$, $C^\xi(T) = 41/15 < 3 = W_p(T)$. For $T \cong T_6(1, 0)$, $C^\xi(T) = 41/12 > 3 = W_p(T)$. For $T \cong P_7$, $C^\xi(T) = 14/5 < 4 = W_p(T)$. For $T \cong T_7(1, 0)$, $C^\xi(T) = 119/60 > 4 = W_p(T)$. For $T \cong T_7(2, 0)$, $C^\xi(T) = 4 = W_p(T)$. For $T \cong T_7(1, 1)$, $C^\xi(T) = 4 = W_p(T)$.

Assume next that $T \not\cong T_n(r, t)$. Then by Lemma 2.7, $W_p(T) \geq n - 2$. Let $P_{d+1} : v_1 v_2 \dots v_d v_{d+1}$ be a diametral path in T . Then

$$C^\xi(G) \leq \frac{2}{4} + \frac{d_2 + d_d}{3} + \frac{2n - 4 - d_2 - d_d}{2} = n - 2 - \frac{d_2 + d_d - 3}{6}$$

where d_2 and d_d are the degrees of the vertices v_2 and v_d , respectively. Thus we have $W_p(T) - C^\xi(T) \geq \frac{d_2 + d_d - 3}{6} > 0$ as $d_2 \geq 2$ and $d_d \geq 2$.

This completes the proof. \square

2.4. Wiener polarity index and average distance

In order to find the relationship between the Wiener polarity index and the average distance, we first consider the following special graphs.

For the complete graph K_n , $\overline{W}(K_n) = 1 > 0 = W_p(K_n)$ for $n \geq 2$.

For the six-membered cycle C_6 , $\overline{W}(C_6) = \frac{9}{5} < 3 = W_p(C_6)$.

For the path P_n , $\overline{W}(P_n) < \frac{n}{2} \leq n - 3 = W_p(P_n)$ for $n \geq 6$.

These examples show that in the general case, WPI and AD are incomparable. Bearing this in mind, we restrict our considerations to trees.

Theorem 2.9. Let T be a tree of order n . If $T \in \{S_n, P_4, P_5\}$, then $\overline{W}(T) \geq W_p(T)$ with equality if and only if $T \cong P_5$. Otherwise,

$$W_p(T) > \overline{W}(T).$$

Proof. We first show that the statement of theorem is true for each tree in the set $\{S_n, P_4, P_5\}$.

If $T \cong S_n$, then $\overline{W}(T) = 2(n-1)/n > 0 = W_p(T)$.

If $T \cong P_4$, then $\overline{W}(T) = 5/3 > 1 = W_p(T)$.

If $T \cong P_5$, then $\overline{W}(T) = 2 = W_p(T)$.

Assume now that $T \notin \{S_n, P_4, P_5\}$. Let d be the diameter of T . Then $d \geq 3$. We consider the following three cases.

Case 1. $3 \leq d \leq n-3$.

By Lemma 2.6, $W_p(T) \geq n-3 \geq d > \overline{W}(T)$.

Case 2. $d = n-2$.

If $T \cong T_n(1, 0)$, then by Lemma 2.6, $W_p(T) = n-3$. As $d \geq 3$, we have $n \geq 5$. If $n = 5$, then $T \cong T_5(1, 0) = S_{1,2}$. Then $W_p(T) = n-3 = 2 > 9/5 = \overline{W}(T)$. So, we may suppose that $n \geq 6$. Then by Lemma 2.1, $\overline{W}(T) \leq \rho < \frac{n}{2}$, and thus,

$$W_p(T) - \overline{W}(T) > n-3 - \frac{n}{2} = \frac{1}{2}(n-6) \geq 0,$$

that is, $W_p(T) > \overline{W}(T)$.

Now, we assume that $T \not\cong T_n(1, 0)$. Then by Lemma 2.6, $W_p(T) \geq n-2 = d > \overline{W}(T)$.

Case 3. $d = n-1$.

Then $T \cong P_n = T_n(0, 0)$. By Lemma 2.6, $W_p(T) = n-3$. Note that $n = d+1 \geq 4$. By our assumption that $T \notin \{S_n, P_4, P_5\}$, we have $n \geq 6$. Thus, $\overline{W}(T) < \rho = \frac{n}{2}$. Therefore, $W_p(T) - \overline{W}(T) > n-3 - \frac{n}{2} = \frac{1}{2}(n-6) \geq 0$.

This completes the proof. \square

References

- [1] Y. Alizadeh, S. Klavžar, Complexity of topological indices: The case of connective eccentric index, MATCH Communications in Mathematical and in Computer Chemistry 76 (2016) 659–667.
- [2] M. Aouchiche, P. Hansen, Proximity and remoteness in graphs: results and conjectures, Networks 58 (2011) 95–102.
- [3] J. A. Bondy, U. S. R. Murty, Graph Theory with Applications, Macmillan, London, 1976.
- [4] F. R. K. Chung, The average distance and the independence number, Journal of Graph Theory 12 (1988) 229–235.
- [5] P. Dankelmann, Average distance and the domination number, Discrete Applied Mathematics 80 (1997) 21–35.
- [6] P. Dankelmann, R. Entringer, Average distance, minimum degree, and spanning trees, Journal of Graph Theory 33 (2000) 1–13.
- [7] K. C. Das, M. Dehmer, Comparison between the zeroth-order Randić index and the sum-connectivity index, Applied Mathematics and Computation 274 (2016) 585–589.
- [8] K. C. Das, I. Gutman, On Wiener and multiplicative Wiener indices of graphs, Discrete Applied Mathematics 206 (2016) 9–14.
- [9] K. C. Das, I. Gutman, M. J. Nadjafi-Arani, Relations between distance-based and degree-based topological indices, Applied Mathematics and Computation 270 (2015) 142–147.
- [10] K. C. Das, M. J. Nadjafi-Arani, Comparison between the Szeged index and the eccentric connectivity index, Discrete Applied Mathematics 186 (2015) 74–86.
- [11] E. de la Viña, B. Waller, Spanning trees with many leaves and average distance, Electronic Journal of Combinatorics 15 (2008) R33.
- [12] A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: theory and applications, Acta Applicanda Mathematicae 66 (2001) 211–249.
- [13] H. Deng, On the extremal Wiener polarity index of chemical trees, MATCH Communications in Mathematical and in Computer Chemistry 66 (2011) 305–314.
- [14] H. Deng, H. Xiao, F. Tang, On the extremal Wiener polarity index of trees with a given diameter, MATCH Communications in Mathematical and in Computer Chemistry 63 (2010) 257–264.
- [15] W. Du, X. Li, Y. Shi, Algorithms and extremal problem on Wiener polarity index, MATCH Communications in Mathematical and in Computer Chemistry 62 (2009) 235–244.
- [16] R. C. Entringer, Distance in graphs: Trees, Journal of Combinatorial Mathematics and Combinatorial Computing 24 (1997) 65–84.
- [17] S. Gupta, M. Singh, A. K. Madan, Connective eccentricity index: a novel topological descriptor for predicting biological activity, Journal of Molecular Graphics and Modeling 18 (2000) 18–25.
- [18] H. Hosoya, Mathematical and chemical analysis of Wiener's polarity number, in: D. H. Rouvray, R.B. King (Eds.), Topology in Chemistry—Discrete Mathematics of Molecules, Horwood, Chichester, 2002, pp. 38–57.

- [19] H. Hua, Y. Chen, K. C. Das, The difference between remoteness and radius of a graph, *Discrete Applied Mathematics* 187 (2015) 103–110.
- [20] H. Hua, K. C. Das, The relationship between the eccentric connectivity index and Zagreb indices, *Discrete Applied Mathematics* 161 (2013) 2480–2491.
- [21] H. Hua, K. C. Das, On the Wiener polarity index of graphs, *Applied Mathematics and Computation* 280 (2016) 162–167.
- [22] O. Ivanciuc, T. S. Balaban, A. T. Balaban, Reciprocal distance matrix, related local vertex invariants and topological indices, *Journal of Mathematical Chemistry* 12 (1993) 309–318.
- [23] B. Liu, H. Hou, Y. Huang, On the Wiener polarity index of trees with maximum degree or given number of leaves, *Computer Mathematics with Applications* 60 (2010) 2053–2057.
- [24] I. Lukovits, W. Linert, Polarity-numbers of cycle-containing structures, *Journal of Chemical Information and Computer Science* 38 (1998) 715–719.
- [25] D. Plavšić, S. Nikolić, N. Trinajstić, Z. Mihalić, On the Harary index for the characterization of chemical graphs, *Journal of Mathematical Chemistry* 12 (1993) 235–250.
- [26] H. Wiener, Structural determination of paraffin boiling points, *Journal of the American Chemical Society* 69 (1947) 17–20.
- [27] K. Xu, M. Liu, K. C. Das, I. Gutman, B. Furtula, A survey on graphs extremal with respect to distance-based topological indices, *MATCH Communications in Mathematical and in Computer Chemistry* 71 (2014) 461–508.
- [28] G. Yu, L. Feng, On the connective eccentricity index of graphs, *MATCH Communications in Mathematical and in Computer Chemistry* 69 (2013) 611–628.
- [29] G. Yu, H. Qu, L. Tang, L. Feng, On the connective eccentricity index of trees and unicyclic graphs with given diameter, *Journal of Mathematical Analysis and Applications* 420 (2014) 1776–1786.
- [30] Y. Zhang, Y. Hu, The Nordhaus–Gaddum-type inequality for the Wiener polarity index, *Applied Mathematics and Computation* 273 (2016) 880–884.